

## ATF-2 (Phospho-Thr73 or 55) Antibody Blocking Peptide

Catalog Number:	51032
Amount:	100µg/100µl
Form of Peptide:	Peptide in 10 mM phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl,
	0.1 mM EDTA, 1 mg/ml BSA, 5% DMF and 5% glycerol.
Peptide Information:	The synthesized phosphopeptide was derived from human ATF-2 around the
	phosphorylation site of threonine 73 or 55 (T-P-T <sup>P</sup> -R-F).
Storage:	Store at -20°C.
Quality Control:	The quality of the peptide was evaluated by reversed-phase HPLC and mass spectrometry.
Specificity:	The peptide specifically blocks the signal of ATF-2 (phospho-Thr73 or 55) antibody (#11032)
	completely in IHC.
Applications:	For Western blotting:add 10 $\mu l$ of antibody and 10 $\mu l$ of blocking peptide to 10 ml of antibody
	dilution buffer, and incubate at 4 $^\circ\!{\rm C}$ over night or at room temperature for 2 hours before
	allowing to react with the blot.
References:	Beausoleil S A, et al. (2004) Proc Natl Acad Sci U S A. 101(33): 12130-12135.
	Abdel-Hafiz H A, et al. (1992) Mol Endocrinol. 6: 2079-2089.
	Gupta S, et al. (1995) Science. 267: 389-393.
	Van Dam H, et al. (1995) EMBO J. 14(8): 1798-1811.
	Livingstone C, et al. (1995) EMBO J. 14: 1785-1797.

