KCNK9 Antibody

Catalog No: #37678

Package Size: #37678-1 50ul #37678-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

KCNK9 Antibody
Rabbit
Polyclonal
Antigen affinity purification.
WB IHC
Hu
The antibody detects endogenous levels of total KCNK9 protein.
Peptide
Synthetic peptide corresponding to a region derived from internal residues of human potassium channel,
subfamily K, member 9
KCNK9
KT3.2; TASK3; K2p9.1; TASK-3
Swiss-Prot#: Q9NPC2NCBI Gene ID: 51305Gene Accssion: NP_001269463
42kd
2.8mg/ml
Rabbit IgG in pH7.3 PBS, 0.05% NaN3, 50% Glycerol.
Store at -20°C

Application Details

Western blotting: 1:500-1:2000 Immunohistochemistry: 1:100-1:300

Images




Gel: 8%SDS-PAGE

Lysates (from left to right): Human paraneoplastic and normal

kidney tissue

Amount of lysate: 40ug per lane Primary antibody: 1/650 dilution Secondary antibody dilution: 1/8000

Exposure time: 40 seconds



Immunohistochemical analysis of paraffin-embedded Human liver cancer tissue using #37678 at dilution 1/40.

Background

This gene encodes a protein that contains multiple transmembrane regions and two pore-forming P domains and functions as a pH-dependent potassium channel. Amplification and overexpression of this gene have been observed in several types of human carcinomas. This gene is imprinted in the brain, with preferential expression from the maternal allele. A mutation in this gene was associated with Birk-Barel mental retardation dysmorphism syndrome. Alternative splicing results in multiple transcript variants.

Published Papers

el at., Potassium channels related to primary aldosteronism: Expression similarities and differences between human and rat adrenals.In Mol Cell Endocrinol on 2015 Dec 5 by Andrew X Chen , Koshiro Nishimoto et al..PMID: 26375812, , (2015)

PMID:26375812

Note: This product is for in vitro research use only and is not intended for use in humans or animals.