

NF-κB p65 Antibody

Catalog No: #48498



Package Size: #48498-1 50ul #48498-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	NF-κB p65 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Peptide affinity purified
Applications	WB, IHC, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	This antibody is produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to N-terminal NF-κB p65.
Other Names	Avian reticuloendotheliosis viral (v rel) oncogene homolog A antibody MGC131774 antibody NF kappa B p65delta3 antibody NFKB3 antibody Nuclear Factor NF Kappa B p65 Subunit antibody Nuclear factor NF-kappa-B p65 subunit antibody Nuclear factor of kappa light polypeptide gene enhancer in B cells 3 antibody Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 antibody OTTHUMP00000233473 antibody OTTHUMP00000233474 antibody OTTHUMP00000233475 antibody OTTHUMP00000233476 antibody OTTHUMP00000233900 antibody p65 antibody p65 NF kappaB antibody p65 NFκB antibody relA antibody TF65_HUMAN antibody Transcription factor p65 antibody v rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B cells 3 (p65)) antibody V rel avian reticuloendotheliosis viral oncogene homolog A antibody v rel reticuloendotheliosis viral oncogene homolog A (avian) antibody V rel reticuloendotheliosis viral oncogene homolog A, nuclear factor of kappa light polypeptide gene enhancer in B cells 3, p65 antibody
Accession No.	Swiss-Prot#:Q04206
Calculated MW	65 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:100-1:200 FC: 1:50-1:100


Images




Western blot analysis of NF-κB p65 on different lysates using anti-NF-κB p65 antibody at 1/1000 dilution. Positive control: Lane1: Hela Lane2: A549 Lane3: PC12 Lane 4: Mouse embryonic stem cell Lane5: NIH/3T3



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-NF-κB p65 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using anti-NF-κB p65 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of Hela cells with NF-κB p65 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Goat anti rabbit IgG (FITC) was used as the secondary antibody.

Background

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. In unstimulated cells, NF-κB is sequestered in the cytoplasm by IκB inhibitory proteins. NF-κB-activating agents can induce the phosphorylation of IκB proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF-κB to enter the nucleus where it regulates gene expression.

References

- 1."Duration of nuclear NF-kappaB action regulated by reversible acetylation." Chen L.F., Fischle W., Verdin E., Greene W.C.Science 293:1653-1657(2001)
- 2."A novel protein overexpressed in hepatoma accelerates export of NF-kappa B from the nucleus and inhibits p53-dependent apoptosis." Higashitsuji H., Higashitsuji H., Nagao T., Nonoguchi K., Fujii S., Itoh K., Fujita J. Cancer Cell 2:335-346(2002)
- 3."Breast cancer metastasis suppressor 1 functions as a corepressor by enhancing histone deacetylase 1-mediated deacetylation of RelA/p65 and promoting apoptosis." Liu Y., Smith P.W., Jones D.R. Mol. Cell. Biol. 26:8683-8696(2006)
- 4."SIRT2 regulates NF-kappaB dependent gene expression through deacetylation of p65 Lys310." Rothgiesser K.M., Erener S., Waibel S., Luscher B., Hottiger M.O. J. Cell Sci. 123:4251-4258(2010)

Published Papers

el at., A multi-omics approach based on 1H-NMR metabonomics combined with target protein analysis to reveal the mechanism of RIAISs on cervical carcinomaIn Aging (Albany NY)On2022 Sep 27byChai Yanlan , Aibibai Jielili et al..PMID:36170024, , (2023)

[PMID:36170024](#)

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF-KB Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy.In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang,et al..PMID: 35370667, , (2022)

[PMID:35370667](#)

el at., Resveratrol attenuates cigarette smoke extract induced cellular senescence in human airway epithelial cells by regulating the miR-34a/SIRT1/NF-kB pathway. In Medicine (Baltimore) on 2022 Nov 18 by Xiao-Li Zeng, Xin-Na Yang, et al..PMID: 36401446, , (2022)

[PMID:36401446](#)

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF-kB Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy. In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang, et al..PMID: 35370667, , (2022)

[PMID:35370667](#)

el at., Salinomycin promotes T-cell proliferation by inhibiting the expression and enzymatic activity of immunosuppressive indoleamine-2,3-dioxygenase in human breast cancer cells. In Toxicol Appl Pharmacol on 2020 Oct 1 by Yuwen Sheng, Zhonghui Zhang, et al..PMID:32822738, , (2020)

[PMID:32822738](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.