

PKR(Phospho-Thr446) Antibody

Catalog No: #11280



Package Size: #11280-1 50ul #11280-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	PKR(Phospho-Thr446) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of PKR only when phosphorylated at threonine 446.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 446 (K-R-T(p)-R-S) derived from Human PKR.
Target Name	PKR
Modification	Phospho
Other Names	ADRB2; E2AK2; EIF2AK2; EIF2aK; PRKR
Accession No.	Swiss-Prot: P19525 NCBI Protein: NP_001129123.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 68kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from K562 cells using PKR(Phospho-Thr446) Antibody #11280(Lane 2) and the same antibody preincubated with blocking peptide(Lane1).

Background

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection.

Abujiang Pataer, et.al. (2002) Cancer Res; 62: 2239.

K. D. Ryman, et.al. (2005) J. Virol; 79: 1487 - 1499.

Susana Guerra, et.al. (2006) J. Biol. Chem; 281: 18734 - 18745.

Published Papers

el at., Fluoxetine Inhibits NLRP3 Inflammasome Activation: Implication in Depression.In Int J Neuropsychopharmacol.On 2016 Sep 21 by Du RH, Tan J et al..PMID:27207922, , (2016)

[PMID:27207922](#)

Suzette Laing, Guohui Wang, Tamara Briazova el at., Airborne Particulate Matter Selectively Activates Endoplasmic Reticulum Stress Response in the Lung and Liver Tissues., Amerian Journal of Physiol Cell Physiol, 299(4):C736-749. doi:10.1152/ajpcell.00529.2009(2010)

[PMID:20554909](#)

el at., Airborne particulate matter selectively activates endoplasmic reticulum stress response in the lung and liver tissues. In Am J Physiol Cell Physiol on 2010 Oct by Suzette Laing, Guohui Wang, et al..PMID: 20554909
, , (2010)

[PMID:20554909](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.