NANOG Antibody

Catalog No: #25045




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$		4.6
	Decri	ption
\boldsymbol{L}	COUL	ווטוו

NANOG Antibody	
Rabbit	
Polyclonal	
Affinity chromatography purified via peptide column	
ELISA WB IHC	
Hu Ms Rt	
Peptide	
Raised against a 19 amino acid peptide near the center of human NANOG.	
NANOG	
NANOG homeobox	
EAW88651	
1mg/ml	
ormulation Supplied in PBS containing 0.02% sodium azide.	
Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of NANOG in human spleen tissue lysate with NANOG antibody at (A) 1 and (B) 2 μ .



Immunohistochemistry of NANOG in human spleen tissue with NANOG antibody at 5 $\mbox{ug/mL}.$

Background

Expression of NANOG is required for the maintenance of pluripotency in epiblast and embryonic stem (ES) cells as well as for the ability to maintain ES self-renewal independently of LIF/Stat3. The role of NANOG in embryonic development suggested that it might be useful in the creation of stem cells that might be useful in cell replacement therapies in the treatment of several degenerative diseases. Artificial stem cells, termed induced pluripotent stem (iPS) cells, can be created by expressing POU5F1 (also known as Oct-4), another germline-specific transcription factor, and the transcription factors Sox2, Klf4 and Lin28 along with c-Myc in mouse fibroblasts. More recently, experiments have demonstrated that iPS cells could be generated using expression plasmids expressing NANOG, Sox2, KlfF4 and c-Myc, eliminating the need for virus introduction, thereby addressing a safety concern for potential use of iPS cells in regenerative medicine.

Published Papers

el at., Wnt10B Is Critical for the Progression of Gastric Cancer.In Oncol Lett on 2017 Jun by Xiao-Dan Wu, Qing-Li Bie,et al..PMID: 28599424, , (2017)
PMID:28599424

Note: This product is for in vitro research use only and is not intended for use in humans or animals.