ERK1/2 (Phospho-Thr202/Tyr204) Antibody

Catalog No: #12082

Package Size: #12082-1 50ul #12082-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


Description	
Product Name	ERK1/2 (Phospho-Thr202/Tyr204) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of ERK1/2 only when phosphorylated at Threonine 202 and Tyrosine
	204 .
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of Threonine 202/Tyrosine 204(F-L-T(p)-E-Y(p)-V-A)Sderived
	from Human ERK1/2.
Target Name	ERK1/2
Modification	Phospho
Other Names	ERK1, ERT2, ERK-1, PRKM3, P44ERK1, ERK1/2
Accession No.	Swiss-Prot#: P27361/P28482; NCBI Gene#: 601795/176948; NCBI Protein#: NP_001035145.1
SDS-PAGE MW	42,44kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C/1 year

Application Details Predicted MW: 42,44kd Western blotting: 1:500~1:1000


Images



Western blot analysis of extracts from 3T3 cells treated with IFNa using ERK1/2 (Phospho-Thr202/Tyr204) Antibody #12082.The lane on the left is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1,ERK 1/2 (phospho Thr202/Y204) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.



Immunofluorescence analysis of mouse-spleen tissue. 1,ERK 1/2 (phospho Thr202/Y204) Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B

Background

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements. The MAPK/ERK cascade plays also a role in initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors. About 160 substrates have already been discovered for ERKs. Many of these substrates are localized in the nucleus, and seem to participate in the regulation of transcription upon stimulation. However, other substrates are found in the cytosol as well as in other cellular organelles, and those are responsible for processes such as translation, mitosis and apoptosis. Moreover, the MAPK/ERK cascade is also involved in the regulation of the endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC); as well as in the fragmentation of the Golgi apparatus during mitosis. The substrates include transcription factors (such as ATF2, BCL6, ELK1, ERF, FOS, HSF4 or SPZ1), cytoskeletal elements (such as CANX, CTTN, GJA1, MAP2, MAPT, PXN, SORBS3 or STMN1), regulators of apoptosis (such as BAD, BTG2, CASP9, DAPK1, IER3, MCL1 or PPARG), regulators of translation (such as EIF4EBP1) and a variety of other signaling-related molecules (like ARHGEF2, FRS2 or GRB10). Protein kinases (such as RAF1, RPS6KA1/RSK1, RPS6KA3/RSK2, RPS6KA2/RSK3, RPS6KA6/RSK4, SYK, MKNK1/MNK1, MKNK2/MNK2, RPS6KA5/MSK1, RPS6KA4/MSK2, MAPKAPK3 or MAPKAPK5) and phosphatases (such as DUSP1, DUSP4, DUSP6 or DUSP16) are other substrates which enable the propagation the MAPK/ERK signal to additional cytosolic and nuclear targets, thereby extending the specificity of the cascade.

Published Papers

el at., Receptor tyrosine kinase C-kit promotes a destructive phenotype of FLS in osteoarthritis via intracellular EMT signalingInMol MedOn2023 Mar 23byXu Cao?1?2,?Song Wu et alPMID:?36959556, , (2023) PMID:36959556
el at., IGF1R-phosphorylated PYCR1 facilitates ELK4 transcriptional activity and sustains tumor growth under hypoxialnNat CommunOn2023 Sep 30byKe Zheng?#?1?2,?Nannan Sha et alPMID:?37777542, , (2023) PMID:37777542
el at., CD155 Cooperates with PD-1/PD-L1 to Promote Proliferation of Esophageal Squamous Cancer Cells via PI3K/Akt and MAPK Signaling Pathways. In Cancers (Basel) on 2022 Nov 15 by Xiyang Tan, Jie Yang, et alPMID:36428703, , (2022) PMID:36428703
el at., Brevilin A Ameliorates Acute Lung Injury and Inflammation Through Inhibition of NF-κB Signaling via Targeting IKKα/β. In Front Pharmacol on 2022 Jun 14 by Lu Liu, Xian Chen,et alPMID:35774606, , (2022) PMID:35774606
el at., CSRP2 suppresses colorectal cancer progression viap130Cas/Racl axis-meditated ERK,PAK, and HIPPOsignaling pathways. In Theranostics on 2020 Sep 2 by Lixia Chen , Xiaoli Long,et al, , (2021) PMID:33042270
el at., CSRP2 suppresses colorectal cancer progression viap130Cas/Racl axis-meditated ERK,PAK, and HIPPOsignaling pathways. In Theranostics on 2020 Sep 2 by Lixia Chen , Xiaoli Long,et alPMID:33042270, , (2021) PMID:33042270
el at., Methamphetamine Blocks Adenosine A2A Receptor Activation via Sigma 1 and Cannabinoid CB1 Receptors. In Int J Mol Sci on 2021 Mar 9 by Mireia Casanovas, Irene
Reyes-Resina, et alPMID:33803075, , (2021) PMID:33803075
el at., Enhancement of anti-acne effect of Scutellaria baicalensis extract by fermentation with symbiotic fungus Penicillium decumbens. In J Biosci Bioeng
on 2020 Nov by Xiaojing Zhu, Yue Mao, et alPMID: 32747300, , (2020) PMID:32747300
el at., The pivotal role of SUMO-1-JNK-Tau axis in an in vitro model of oxidative stress counteracted by the protective effect of curcumin. In Biochem Pharmacol on 2020 Aug by Lucia Buccarello, Jessica Dragotto, et alPMID:32502496, , (2020) PMID:32502496
el at., Glutamate Attenuates the Survival Property of IGFR through NR2B Containing N-Methyl-D-aspartate Receptors in Cortical Neurons. In Oxid Med Cell Longev on 2020 Aug 11 by Xia Zhao, Chao Han, et alPMID:32849999, , (2020) PMID:32849999
el at., Activation of GCN2/ATF4 signals in amygdalar PKC-δ neurons promotes WAT browning under leucine deprivation. In Nat Commun on 2020 Jun 5 by Feixiang Yuan, Haizhou Jiang, et alPMID:32504036, , (2020) PMID:32504036
el at., Carbon monoxide Induces the assembly of stress granule through the Integrated stress response. In Biochem Biophys Res Commun on 2019 Apr 30 by Chen Y, Joe Y, et alPMID:30885431, , (2019) PMID:30885431

el at., miR-141-5p regulate ATF2 via effecting MAPK1/ERK2 signalIng to promote preeclampsia.Biomed Pharmacother. In Biomed Pharmacother on 2019 May 7 by Wang Y1, Cheng K, et al..PMID: 31075732, , (2019)

PMID:31075732

el at., HMGN2 regulates non-tuberculous mycobacteria survival via modulation of M1 macrophage polarization. In J Cell Mol Med on 2019 Dec by Wang X, Chen S, et al. PMID:31596045, , (2019)

PMID:31596045

el at., The synergic Inhibitory effects of dark tea (Camellia sInensis) extract and p38 Inhibition on the growth of pancreatic cancer cells. In J Cancer on 2019 Oct 21 by Zheng K, Zhao Q, et al. PMID:31777585, , (2019)

PMID:31777585

el at., The protective effects of β-sitosterol and vermicularIn from Thamnolia vermicularis (Sw.) Ach. agaInst skIn agIng In vitro. In An Acad Bras Cienc on 2019 Dec 2 by Haiyuan YU, Shen X,et al..PMID:31800700, , (2019)

PMID:31800700

el at., Astragalus polysaccharides exerts anti-infective activity by inducing human cathelicidin antimicrobial peptide LL-37 in respiratory epithelial cells.In Phytother Res. On 2018 Aug by Zhao L, Tan S et al.. PMID: 29672953, , (2018)

PMID:29672953

el at., Crucial Roles of 5-HT and 5-HT2 Receptor in Diabetes-Related Lipid Accumulation and Pro-Inflammatory Cytokine Generation in Hepatocytes.In Cell Physiol Biochem.On 2018 Aug 17 by Fu J, Li C et al..PMID: 30121645, , (2018) PMID:30121645

el at., Inhibition of TPL2 by interferon-α suppresses bladder cancer through activation of PDE4D.In J Exp Clin Cancer Res. On 2018 Nov 27 by Qiang Z, Zhou ZY et al.. PMID: 30482227, , (2018)

PMID:30482227

el at., Ajudecumin A from Ajuga ovalifolia var. calantha exhibits anti-inflammatory activity in lipopolysaccharide-activated RAW264.7 murine macrophages and animal models of acute inflammation. In Pharm Biol. On 2018 Dec by Zhang H, Ren QC et al.. PMID: 31070535 , , (2018) PMID:31070535

el at., Phosphoglycerate Kinase 1 Phosphorylates Beclin1 to Induce Autophagy.In Mol Cell on 2017 Mar 2 by Xu Qian, Xinjian Li, et al.. PMID: 28238651, , (2017)

PMID:28238651

el at., Salidroside pretreatment attenuates apoptosis and autophagy during hepatic ischemial ζ • eperfusion injury by inhibiting the mitogen-activated protein kinase pathway in mice. In Drug Des Devel Ther on 2017 Jul 3 by Jiao Feng, Qinghui Zhang, et al.. PMID: 28721018, , (2017) PMID:28721018

el at., Puerarin Inhibits Expression of Tissue Factor Induced by Oxidative Low-Density Lipoprotein Through Activating the PI3K/Akt/eNOS Pathway and Inhibiting Activation of ERK1/2 and NF-I-• B.In Life Sci on 2017 Dec 15 by Hua-Fei Deng, Xiao-Li Wang, et al.. PMID: 29037842, , (2017) PMID:29037842

el at., Capsular polysaccharide of mycoplasma ovipneumoniae induces sheep airway epithelial cell apoptosis via ROS-dependent JNK/P38 MAPK pathways. In Oxid Med Cell Longev on 2017 Mar 7 by Zhongjia Jiang, Fuyang Song, et al.. PMID: 28367270, , (2017) PMID:28367270

el at., Ellagic Acid Inhibits RANKL-induced Osteoclast Differentiation by Suppressing the p38 MAP Kinase Pathway. In Arch Pharm Res on 2017 Jan by Mpho Rantlha, Travers Sagar, et al.. PMID: 27384064, , (2017)

PMID:27384064

el at., Functional ζE--Opioid-Galanin Receptor Heteromers in the Ventral Tegmental Area.In J Neurosci on 2017 Feb 1 by Estefan a Moreno. Cθ

sar Quiroz, et al.. PMID: 28007761, , (2017)

PMID:28007761

el at., Altered Timing of Riboflavin and Ultraviolet Light Pathogen Inactivation Improves Platelet in Vitro Quality. In Transfusion on 2017 Aug by Peter Schubert, Brankica Culibrk, et al.. PMID: 28500654, , (2017)

PMID:28500654

el at., Neurokinin B Exerts Direct Effects on the Ovary to Stimulate Estradiol Production. In Endocrinology. On 2016 Sep by Qi X, Salem M et

al..PMID:27580802, , (2016)

PMID:27580802

el at., FSTL1 as a Potential Mediator of Exercise-Induced Cardioprotection in Post-Myocardial Infarction Rats. In Sci Rep on 2016 Aug 26 by Yue Xi, Da-Wei Gong et al.. PMID: 27561749, (2016)

PMID:27561749

el at., Response of MAPK pathway to iron oxide nanoparticles in vitro treatment promotes osteogenic differentiation of hBMSCs. In Biomaterials. On 2016 Apr by Wang Q, Chen B et al.. PMID: 26874888, (2016)

PMID:26874888

el at., Bruceine D induces apoptosis in human chronic myeloid leukemia K562 cells via mitochondrial pathway. In Am J Cancer Res on 2016 Mar 15

by Jian-Ye Zhang, Min-Ting Lin et al.. PMID: 27186433, , (2016)

PMID:27186433

el at., An alternatively spliced variant of CXCR3 mediates the metastasis of CD133+ liver cancer cells induced by CXCL9.In Oncotarget on 2016 Mar 22 by Qiang Ding, Yujia Xia et al..PMID:26883105, (2016)

PMID:26883105

el at., Genipin derivatives protect RGC-5 from sodium nitroprusside-induced nitrosative stress. In Int J Mol Sci on 2016 Jan 19 by Rikang Wang, Jiaqiang Zhao, et al..PMID:26797604

, , (2016)

PMID:26797604

el at., Mycoplasma ovipneumoniae induces sheep airway epithelial cell apoptosis through an ERK signalling-mediated mitochondria pathway. In BMC Microbiol on 2016 Sep 23 by Yanan Li, Zhongjia Jiang, et al.. PMID: 27663303, , (2016)

PMID:27663303

el at., Inhibitory effects of eugenol on RANKL-induced osteoclast formation via attenuation of NF-I-• B and MAPK pathways.In Connect Tissue Res on 2015 Jun by Vishwa Deepak, Abe Kasonga et al..PMID:25405641, (2015)

PMID:25405641

el at., Overexpression of astrocyte-elevated gene-1 is associated with ovarian cancer development and progression. In Mol Med Rep on 2015 Apr by Bo Zhou, Jue Yang et al..PMID:25483832, (2015)

PMID:25483832

el at., Combinational treatment of curcumin and quercetin against gastric cancer MGC-803 cells in vitro Molecules on 2015 Jun 22 by Jian-Ye Zhang , Min-Ting Lin et al..PMID:26111180 , , (2015)

PMID:26111180

el at., CPU-12, a novel synthesized oxazolo [5, 4-d] pyrimidine derivative, showed superior anti-angiogenic activity. In J Pharmacol Sci on 2015 Sep by Jiping Liu, Ya-Hui Deng et al.. PMID: 26154846, , (2015)

PMID:26154846

el at., Piperine alleviates osteoclast formation through the p38/ci ζ os/NFATc1 signaling axis.In Biofactors on Nov-Dec 2015 by Vishwa Deepak , Marlena C Kruger et al..PMID: 26627060, , (2015)

PMID:26627060

el at., Exosomes Derived From Human Mesenchymal Stem Cells Confer Drug Resistance in Gastric Cancer. In Cell Cycle on 2015 Aug 3 by Runbi Ji, Bin Zhang et al., PMID:26091251, , (2015)

PMID:26091251

el at., Cannabinoid receptor agonist WIN55,212-2 and fatty acid amide hydrolase inhibitor URB597 suppress chronic cerebral hypoperfusion-induced neuronal apoptosis by inhibiting c-Jun N-terminal kinase signaling.In Neuroscience 2015 Aug 20 by S-H Su , Y-F Wu et al..PMID:25795598, , (2015) PMID:25795598

el at., Role of SMC1A overexpression as a predictor of poor prognosis in late stage colorectal cancer. In BMC Cancer on 2015 Mar 4 by Jianwei

Wang , Shaojun Yu et al.. PMID: 25884313, , (2015)

PMID:25884313

el at., Neutral phosphate-affinity SDS-PAGE system for profiling of protein phosphorylation. In Methods Mol Biol on 2015 by Eiji Kinoshita, Tohru

Koikeı ε t al..PMID:25820732

, , (2015)

PMID:25820732

el at., Celecoxib ameliorates portal hypertension of the cirrhotic rats through the dual inhibitory effects on the intrahepatic fibrosis

and angiogenesis. In PLoS One on 2013 Jul 26 by Gao JH, Wen SL, et al.. PMID: 23922700, , (2013)

PMID:23922700

el at., Knockdown of Protein Phosphatase Magnesium-Dependent 1 (PPM1D) Through Lentivirus-Mediated RNA Silencing Inhibits Colorectal

Carcinoma

Cell Proliferation. In Technol Cancer Res Treat on 2013 Dec by Hongzhuan Yin, Zhaopeng Yan, et al.. PMID: 23745790, , (2013)

PMID:23745790

el at., Asperosaponin VI, a saponin component from Dipsacus asper wall, induces osteoblast differentiation through bone morphogenetic

protein-2/p38 and extracellular signal-regulated kinase 1/2 pathway. In Phytother Res on 2011 Nov by Niu Y, Li Y, et al. PMID: 21452371, , (2011) PMID: 21452371

el at., The pivotal role of SUMO-1-JNK-Tau axis in an in vitro model of oxidative stress counteracted by the protective effect of curcumin. In Biochem Pharmacol on 2020 Aug by Lucia Buccarello, Jessica Dragotto, et al..PMID:32502496, , ()

PMID:32502496

Note: This product is for in vitro research use only and is not intended for use in humans or animals.