GSK3β(Phospho-Ser9) Antibody

Catalog No: #11002

Package Size: #11002-1 50ul #11002-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description	

Product Name	GSK3β(Phospho-Ser9) Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.		
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho		
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.		
Applications	WB IHC IF		
Species Reactivity	Hu Ms Rt		
Specificity	The antibody detects endogenous level of GSK3 beta only when phosphorylated at serine 9.		
Immunogen Type	Peptide-KLH		
Immunogen Description	Synthesized phospho-peptide around the phosphorylation site of human GSK3 β (phospho Ser9)		
Target Name	GSK3β		
Modification	Phospho		
Other Names	Factor A, GSK-3 beta, Protein kinase GSK-3-beta, kinase GSK-3 beta		
Accession No.	Swiss-Prot: P49841NCBI Protein: NP_x.1		
SDS-PAGE MW	46kd		
Concentration	1.0mg/ml		
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Storage	Store at -20°C for long term preservation (recommended). Avoid Store at 4°C for short term use.		


Application Details

Predicted MW: 46kd		
Western blotting: 1:500~1:1000		
Immunohistochemistry: 1:50~1:100		
Immunofluorescence: 1:100~1:200		

Images




Western blot analysis of extracts from 293 cells untreated(lane 1) or treated with serum(lane 2) using GSK3 β (Phospho-Ser9) Antibody #11002.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using GSK3β (Phospho-Ser9) Antibody #11002 (left) or the same antibody preincubated with blocking peptide #51002 (right).




Immunofluorescence staining of methanol-fixed Hela cells showing cytoplasmic staining using GSK3β(Phospho-Ser9) Antibody #11002.



Immunohistochemical analysis of paraffin-embedded human Lung carcinoma tissue using GSK3β (Phospho-Ser9) Antibody #11002.



Immunofluorescence staining of methanol-fixed Hela cells showing cytoplasmic staining using GSK3 β (Phospho-Ser9) Antibody #11002.



Western blot analysis of extracts from Hela cells, treated with EGF or calf intestinal phosphatase (CIP), using GSK3 β (Phospho-Ser9) Antibody #11002.

Background

Participates in the Wnt signaling pathway. Implicated in the hormonal control of several regulatory proteins including glycogen synthase, MYB and the transcription factor JUN. Phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA. Phosphorylates MUC1 in breast cancer cells, and decreases the interaction of MUC1 with CTNNB1/beta-catenin. Phosphorylates CTNNB1/beta-catenin. Fan G, et al. (2003) J Biol Chem. 278(52): 52432-52436. Barry FA, et al. (2003) FEBS Lett. 553(1-2): 173-178. Welsh, et al. (1996) Trends Cell Biol. 6: 274-279. Srivastava A K, et al. (1998) Mol Cell Biochem. 182: 135-141.

Published Papers

el at., Periostin secreted by cancer-associated fibroblasts promotes cancer stemness in head and neck cancer by activating protein tyrosine kinase 7.In Cell Death Dis.On 2018 Oct 22 by Yu B, Wu K et al..PMID: 30348980, , (2018)

PMID:30348980

el at., Epithelial Cells Attenuate Toll-Like Receptor-Mediated Inflammatory Responses in Monocyte-Derived Macrophage-Like Cells to Mycobacterium tuberculosis by Modulating the PI3K/Akt/mTOR Signaling Pathway. In Mediators Inflamm. On 2018 Sep by Yang Y, Sun Y et al.. PMID:30356420, ,

(2018)

PMID:30356420

el at., Gastrokine-2 suppresses epithelial mesenchymal transition through PI3K/AKT/GSK3ε[°]Y signaling in gastric cancer.In Tumour Biol.On 2016 Sep by Dai J, Qian C et al..PMID:27323966, , (2016)

PMID:27323966

el at., Overexpression of astrocyte-elevated gene-1 is associated with ovarian cancer development and progression. In Mol Med Rep on 2015 Apr by Bo Zhou, Jue Yang et al..PMID:25483832, (2015)

PMID:25483832

el at., Fumosorinone, a Novel PTP1B Inhibitor, Activates Insulin Signaling in Insulin-Resistance HepG2 Cells and Shows Anti-Diabetic Effect in Diabetic KKAy Mice .In Toxicol Appl Pharmacol On 2015 May 15 by Zhi-Qin Liu, Ting Liu et al..PMID: 25796170

, , (2015)

PMID:25796170

el at., SNAILs promote G1 phase in selected cancer cells.In Int J Oncol on 2015 Nov by Ya-Lan Wu, Jian-Xin Xue et al ...PMID:26352801, , (2015) PMID:26352801

el at., A ε[°]Y1/2 Adrenergic Receptori ζ ±ensitive Intracellular Signaling Pathway Modulates CCL2 Production in Cultured Spinal Astrocytes.In J Cell Physiol on 2014 Mar by Norimitsu Morioka, Hiromi Abe et al..PMID:24037783, , (2014)

PMID:24037783

Estefanq— I a de Munck, Emma Mu?noz-Sq— ϵ ez, Bego?na G. Miguel el at., B-N-methylamino-l-alanine causes neurological andpathological phenotypes mimicking Amyotrophic LateralSclerosis (ALS): The first step towards an experimentalmodel for sporadic ALS., environmental toxicology and pharmacology, 36:243q— C255(2013)

PMID:23688553

Yu-fei Pan, Li-wei Dong, Min Wang el at., Signal regulatory protein ε° θ negatively regulates mast-cell activation following Fc ε° θ RI aggregation., Eur. J. Immunol., 43(6):1598n— C1607(2013)

PMID:23504624

el at., Phosphoproteome analysis reveals an important role for glycogen synthase kinase-3 in perfluorododecanoic acid-induced rat liver toxicity. In Toxicol Lett on 2013 Mar 27 by Hongxia Zhang, Junjie Hou, et al..PMID: 23353032, (2013)

PMID:23353032

el at., ε[°]Y-N-methylamino-l-alanine causes neurological and pathological phenotypes mimicking Amyotrophic Lateral Sclerosis (ALS): The first step towards an experimental model for sporadic ALS. In Environ Toxicol Pharmacol on 2013 Sep by Estefanı a de Munck, Emma MuέE½oz-Sθ°©ez, et al..PMID: 23688553, , (2013)

PMID:23688553

el at., Signal regulatory protein δO negatively regulates masti ζ • ell activation following Fc θ RI aggregation. In Eur J Immunol on 2013 Jun by Yu-fei Pan, Li-wei Dong, et al..PMID: 23504624, , (2013)

PMID:23504624

el at., Quantitative proteomics reveals that miR-155 regulates the PI3K-AKT pathway in diffuse large B-cell lymphoma. In Am J Pathol on 2012 Jul by Huang X, Shen Y, et al..PMID:22609116, , (2012)

PMID:22609116

Yingjuan Yang, Jinzeng Yang, Rongxin Liu el at., Accumulation of ε θ -catenin by lithium chloride in porcine myoblast cultures accelerates cell differentiation., Molecular Biology Reports, Volume 38, Number 3, Pages 2043-2049(2011)

PMID:20857211

el at., Accumulation of ɛ°Y-Catenin by Lithium Chloride in Porcine Myoblast Cultures Accelerates Cell Differentiation. In Mol Biol Rep on 2011 Mar by

Yingjuan Yang, Jinzeng Yang, et al.. PMID: 20857211, , (2011)

PMID:20857211

Cong REN, Jia-Mou LI, Xin LIN el at., LIPUS Enhance Elongation of Neurites in Rat Cortical Neurons through Inhibition of GSK-3 $\epsilon^{"}$ θ ., Biomedical and Environmental Sciences, 23(3):244-249(2010)

PMID:20708505

Wen-Fei Tan, Xue-Zhao Cao, Jun-KeWang el at., Protective effects of lithium treatment for spatial memory deficits induced by tau hyperphosphorylation in splenectomized rats., Clinical and Experimental Pharmacology and Physiology., 37(10):1010n— C1015(2010) PMID:20659131

el at., Protective effects of lithium treatment for spatial memory deficits induced by tau hyperphosphorylation in splenectomized rats.In

Clin Exp Pharmacol Physiol on 2010 Oct by Tan WF, Cao XZ,et al..PMID:20659131, , (2010)

PMID:20659131

el at., LIPUS Enhance Elongation of Neurites in Rat Cortical Neurons Through Inhibition of GSK-3beta. In Biomed Environ Sci on 2010 Jun by Cong Ren, Jia-Mou Li, et al..PMID:20708505, , (2010)

PMID:20708505

Note: This product is for in vitro research use only and is not intended for use in humans or animals.