

## Nanog Antibody

Catalog No: #21423

Package Size: #21423-1 50ul #21423-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description


Product Name	Nanog Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous level of total Nanog protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.137~141 (K-Q-V-K-T) derived from Nano g
Target Name	Nanog
Other Names	Homeobox transcription factor Nanog
Accession No.	Swiss-Prot: Q9H9S0NCBI Protein: NP_079141.2
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

## Application Details

Predicted MW: 42kd

Western blotting: 1:1000

## Images



Western blot analysis of extracts from human Umbilical cord mesenchymal stem cell using Nano g Antibody #21423.

## Background

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and

prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes By similarity. Acts as a transcriptional activator or repressor By similarity. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3' By similarity. When overexpressed, promotes cells to enter into S phase and proliferation Do HJ, et al.Biochem Biophys Res Commun. 2007 Feb 16;353(3):770-5.  
Boyer LA, et al.Cell. 2005 Sep 23;122(6):947-56.  
Freberg CT, et al. Mol Biol Cell. 2007 May;18(5):1543-53.  
Chambers I, et al. Cell. 2003 May 30;113(5):643-55.

## Published Papers

el at., SALL4 promotes gastric cancer progression through activating CD44 expression.In Oncogenesis on 2016 Nov 7 by X Yuan , X Zhang et al..PMID:27819668 , , (2016)

[PMID:27819668](#)

el at., HucMSC exosome-delivered 14-3-3 $\zeta$  orchestrates self-control of the Wnt response via modulation of YAP during cutaneous regeneration. In Stem Cells on 2016 Oct by Bin Zhang, Yinghong Shi, et al..PMID: 27334574, , (2016)

[PMID:27334574](#)

el at., Tumorigenic hybrids between mesenchymal stem cells and gastric cancer cells enhanced cancer proliferation, migration and stemness.In BMC Cancer on 2015 Oct 24 by Jianguo Xue , Yuan Zhu et al..PMID: 26498753, , (2015)

[PMID:26498753](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.