

STAT1(Phospho-Tyr701) Antibody

Catalog No: #11044



Package Size: #11044-1 50ul #11044-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	STAT1(Phospho-Tyr701) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of STAT1 only when phosphorylated at tyrosine 701.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 701 (T-G-Y(p)-I-K) derived from Human STAT1.
Target Name	STAT1
Modification	Phospho
Other Names	CANDF7, ISGF-3, STAT91
Accession No.	Swiss-Prot: P42224NCBI Protein: NP_009330.1
SDS-PAGE MW	84,91kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 84,91kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100


Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using STAT1(Phospho-Tyr701) Antibody #11044(left) or the same antibody preincubated with blocking peptide(right).



Western blot analysis of extracts from MEF cells untreated or treated with interferon- γ (IFN γ) using STAT1 (Phospho-Tyr701) Antibody #11044.



Western blot analysis of extracts from HeLa cells, treated with IFN α or calf intestinal phosphatase (CIP), using STAT1 (Phospho-Tyr701) Antibody #11044.

Background

Signal transducer and activator of transcription that mediates signaling by interferons (IFNs). Following type I IFN (IFN- α and IFN- β) binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. In response to type II IFN (IFN- γ), STAT1 is tyrosine- and serine-phosphorylated. It then forms a homodimer termed IFN- γ -activated factor (GAF), migrates into the nucleus and binds to the IFN γ activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state.

Heim M H, (1999) J Recept Signal Transduct Res. 19: 75-120.

Durbin J E, et al. (1996) Cell. 84: 443-450.

Meraz M A, et al. (1996) Cell. 84: 431-442.

Wakao H, et al. (1994) EMBO J. 13: 2182-2191.

Demoulin J, B. et al. (1999) J Biol Chem. 274: 25855-258

Ihle J N, et al. (1994) Trends Biochem Sci. 19: 222-227.

Published Papers

el at., Costunolide improved dextran sulfate sodium-induced acute ulcerative colitis in mice through NF- κ B, STAT1/3, and Akt signaling pathways. In Int Immunopharmacol on 2020 Jul by Fan Xie, Hai Zhang, et al..PMID:32413737, , (2020)

[PMID:32413737](#)

el at., Chronic Inflammation In Response to Injury: Retention of Myeloid Cells in Injured Tissue Is Driven by Myeloid Cell IntrInsic Factors. In J Invest Dermatol on 2019 Jan 28 by Torbica T, Wicks K, et al..PMID:30703358, , (2019)

[PMID:30703358](#)

el at., A thiazole-derived oridonIn analogue exhibits antitumor activity by directly and allosterically InhibitIng STAT3. In J Biol Chem on 2019 Nov 15;by Shen X, Zhao L,et al..PMID:31594861, , (2019)

[PMID:31594861](#)

el at., Danzhi Jiangtang Capsule ameliorates kidney injury via inhibition of the JAK-STAT signaling pathway and increased antioxidant capacity in

STZ-induced diabetic nephropathy rats. In Biosci Trends. On 2019 Jan 22 by Sun M, Bu W et al.. PMID: 30606979 , , (2018)

[PMID:30606979](#)

el at., DJC suppresses Advanced Glycation End Products-Induced JAK-STAT signaling and ROS in mesangial cells. In Evid Based Complement Alternat Med on 2017 May 29 by Min Sun, Yan Li ,et al.. PMID: 28630633, , (2017)

[PMID:28630633](#)

el at., Inhibitory Effects of Chalcone Glycosides Isolated From Brassica Rapa L. 'Hidabeni' and Their Synthetic Derivatives on LPS-induced NO Production in Microglia. In Bioorg Med Chem on 2011 Sep 15 by Hirokazu Hara, Yoko Nakamura, et al.. PMID: 21856162, , (2017)

[PMID:21856162](#)

el at., Expression of interferon effector gene SART1 correlates with interferon treatment response against hepatitis B infection. In Mediators Inflamm on 2016 by Yong Li, Chuanlong Zhu et al.. PMID: 28077916, , (2016)

[PMID:28077916](#)

el at., Mycobacterium tuberculosis EspB protein suppresses interferon- γ -induced autophagy in murine macrophages. In J Microbiol Immunol Infect on 2016 Dec by DanDan Huang, Lang Bao et al.. PMID:25641595 , , (2016)

[PMID:25641595](#)

el at., Newly Synthesized α - ζ -hidabeni- β -Chalcone Derivatives Potently Suppress LPS-Induced NO Production via Inhibition of STAT1, but Not NF- κ B, JNK, and p38, Pathways in Microglia. In Biol Pharm Bull On 2014 by Hirokazu Hara , Ryoko Ikeda et al.. PMID: 24882415, , (2014)

[PMID:24882415](#)

el at., Molecular analysis of curcumin-induced polarization of murine RAW264.7 macrophages. In J Cardiovasc Pharmacol on 2014 Jun by Fangyuan Chen , Ning Guo et al.. PMID:24709638, , (2014)

[PMID:24709638](#)

el at., The C-terminal Domain of Chikungunya Virus nsP2 Independently Governs Viral RNA Replication, Cytopathicity, and Inhibition of Interferon Signaling. In J Virol on 2013 Sep by Jelke J Fros, Erika van der Maten, et al.. PMID:23864632, , (2013)

[PMID:23864632](#)

C.-C.E. Lan, C.-S. Wu, S.-M. Huang et al., High-glucose environment reduces human α -defensin-2 expression in human keratinocytes: implications for poor diabetic wound healing., British Association of Dermatologists, 166(6):1221-1229 (2012)

[PMID:22283836](#)

el at., High-glucose Environment Reduces Human α -Defensin-2 Expression in Human Keratinocytes: Implications for Poor Diabetic Wound Healing. In

Br J Dermatol on 2012 Jun by C-C E Lan, C-S Wu, et al.. PMID:22283836, , (2012)

[PMID:22283836](#)

Hirokazu Hara, Yoko Nakamura, Masayuki Ninomiya et al., Inhibitory effects of chalcone glycosides isolated from Brassica rapa L.

α -hidabeni- β and their synthetic derivatives on LPS-induced NO production in microglia., Bioorganic & Medicinal Chemistry, 19(18):5559-5568(2011)

[PMID:21856162](#)

el at., Inhibitory effects of chalcone glycosides isolated from Brassica rapa L. α -hidabeni- β and their synthetic derivatives on LPS-induced NO production in microglia. In Bioorg Med Chem on 2011 Sep 15 by Hirokazu Hara, Yoko Nakamura, et al.. PMID:21856162, , (2011)

[PMID:21856162](#)

Hsin-Chien Chen, Hsin-I Ma, Huey-Kang Sytwu et al., Neural Stem Cells Secrete Factors That Promote Auditory Cell Proliferation Via a Leukemia Inhibitory Factor Signaling Pathway, Journal of Neuroscience Research, 88:3308-3318 (2010)

[PMID:20882565](#)

Jelke J. Fros, Wen Jun Liu, Natalie A. Prow et al., Chikungunya virus nonstructural protein 2 inhibits type I/II interferon-stimulated JAK-STAT signaling., Journal of Virology, 84(20):10877-10887(2010)

[PMID:20686047](#)

et al., Neural stem cells secrete factors that promote auditory cell proliferation via a leukemia inhibitory factor signaling pathway. In J Neurosci Res on 2010 Nov 15 by Hsin-Chien Chen, Hsin-I Ma, et al..PMID: 20882565, , (2010)

[PMID:20882565](#)

et al., Chikungunya virus nonstructural protein 2 inhibits type I/II interferon-stimulated JAK-STAT signaling. In J Virol on 2010 Oct by Jelke J Fros, Wen Jun Liu, et al..PMID:

20686047, , (2010)

[PMID:20686047](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.