

NFkB-p65(Phospho-Ser536) Antibody

Catalog No: #11014



Package Size: #11014-1 50ul #11014-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	NFkB-p65(Phospho-Ser536) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC IF
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of NF-kB p65 only when phosphorylated at serine 536.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 536 (F-S-S(p)-I-A) derived from Human NFkB-p65.
Conjugates	Unconjugated
Target Name	NFkB-p65
Modification	Phospho
Other Names	NFKB3; RELA; TF65; Transcription factor p65; p65
Accession No.	Swiss-Prot: Q04206NCBI Protein: NP_001138610.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 65kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images




Western blot analysis of extracts from HepG2 and HL60 cells untreated or treated with TNF using NFκB-p65(Phospho-Ser536) Antibody #11014.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using NFκB-p65 (Phospho-Ser536) Antibody #11014 (left) or the same antibody preincubated with blocking peptide #51014 (right).




Immunofluorescence staining of methanol-fixed HeLa cells using NFκB-p65(Phospho-Ser536) Antibody #11014.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using NFκB-p65 (Phospho-Ser536) Antibody #11014.



Immunohistochemical analysis of paraffin-embedded human Lung carcinoma tissue using NFκB-p65 (Phospho-Ser536) Antibody #11014.



Immunofluorescence staining of methanol-fixed MEF cells using NFκB-p65 (Phospho-Ser536) Antibody #11014.

Background

NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-kappa-B p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of I-kappa-B upon NF-kappa-B in the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex.

Doyle S L, et al. (2005) J Biol Chem. 280(25): 23496-23501.

Anwar K N, et al. (2004) J Immunol. 173(11): 6965-6972.

Baeuerle P A, et al. (1994) Annu Rev Immunol. 12:141-179.

Baeuerle P A, et al. (1996) Cell 87:13-20.

Haskill S, et al. (1991) Cell 65:1281-1289.

Published Papers

Jitakshi De, Robert E. Brown et al., Tissue-microarray based immunohistochemical analysis of survival pathways in nodular sclerosing classical Hodgkin lymphoma as compared with Non-Hodgkin's lymphoma. , Int J Clin Exp Med, 3(1):55-68(2010)

[PMID:20369041](#)

et al., Downregulated Expression of Solute Carrier Family 26 Member 6 in NRK-52E Cells Attenuates Oxalate-Induced Intracellular Oxidative Stress. In Oxid Med Cell Longev. On 2018 Oct 10 by Jiang H, Gao X et al.. PMID:30405874, , (2018)

[PMID:30405874](#)

et al., Advanced Glycation End Products Stimulate Angiotensinogen Production in Renal Proximal Tubular Cells. In Am J Med Sci on 2019 Jan by Garagliano JM, Katsurada A, et al.. PMID:30466736, , (2019)

[PMID:30466736](#)

et al., Activation of α7-nAChRs protects SH-SY5Y cells from 1-methyl-4-phenylpyridinium-Induced apoptotic cell death via ERK/p53 signalling pathway. In J Cell Physiol on 2019 Mar 25 by Xu S, Yang B, et al.. PMID:30912145, , (2019)

[PMID:30912145](#)

et al., Nuclear factor I A promotes temozolomide resistance in glioblastoma via activation of nuclear factor κB pathway. In Life Sci on 2019 Nov 1 by Yu X, Wang M, et al.. PMID:31614149, , (2019)

[PMID:31614149](#)

el at., CYP4A11 is involved in the development of nonalcoholic fatty liver disease via ROS-induced lipid peroxidation and inflammation. In *Int J Mol Med* on 2020 Apr by Gao H, Cao Y, et al..PMID:32124935, , (2020)

[PMID:32124935](#)

el at., Induction of apoptosis through extrinsic/intrinsic pathways and suppression of ERK/NF- κ B signalling participate in anti-glioblastoma of imipramine. In *J Cell Mol Med* on 2020 Apr; by Hsu FT, Chiang IT et al..PMID:32149465, , (2020)

[PMID:32149465](#)

el at., Phytoestrogens protect joints in collagen induced arthritis by increasing IgG glycosylation and reducing osteoclast activation. In *Int Immunopharmacol* on 2020 Mar 12 by Du N, Song L, et al..PMID:32172207, , (2020)

[PMID:32172207](#)

el at., Blockade of extracellular high-mobility group box 1 attenuates inflammation-mediated damage and haze grade in mice with corneal wounds. In *Int Immunopharmacol* on 2020 Apr 9; by Zhou Y, Wang T, et al..PMID:32279044, , (2020)

[PMID:32279044](#)

el at., Tissue-microarray Based Immunohistochemical Analysis of Survival Pathways in Nodular Sclerosing Classical Hodgkin Lymphoma as Compared With Non-Hodgkin's Lymphoma. In *Int J Clin Exp Med* on 2010 Jan 30 by Jitakshi De, Robert E Brown et al..PMID:20369041, , (2010)

[PMID:20369041](#)

el at., Dietary Oxidized n-3 PUFA Induce Oxidative Stress and Inflammation: Role of Intestinal Absorption of 4-HHE and Reactivity in Intestinal Cells. In *J Lipid Res* on 2012 Oct by Manar Awada, Christophe O Soulage, et al..PMID:22865918, , (2012)

[PMID:22865918](#)

el at., Ent-11 δ O⁻-hydroxy-15-oxo-kaur-16-en-19-oic-acid induces apoptosis and cell cycle arrest in CNE-2Z nasopharyngeal carcinoma cells. In *Oncol Rep* on 2013 Jun by Kefeng Wu, Yi Liu, et al..PMID: 23563985, , (2013)

[PMID:23563985](#)

el at., NDR1/STK38 potentiates NF- κ B activation by its kinase activity. In *Cell Biochem Funct* on 2012 Dec by Hu Shi, Dan Lu, et al..PMID: 22674419, , (2012)

[PMID:22674419](#)

el at., Pterisolic acid G, a novel ent- ζ -aurane diterpenoid, inhibits viability and induces apoptosis in human colorectal carcinoma cells. In *Oncol Lett* on 2017 Nov by Shuangli Qiu, Xin Wu, et al..PMID: 29113182, , (2017)

[PMID:29113182](#)

el at., Capsular Polysaccharide Is a Main Component of *Mycoplasma ovipneumoniae* in the Pathogen-Induced Toll-Like Receptor-Mediated Inflammatory Responses in Sheep Airway Epithelial Cells. In *Mediators Inflamm* on 2017 May 3 by Zhongjia Jiang, Fuyang Song, et al..PMID: 28553017, , (2017)

[PMID:28553017](#)

el at., Pre-treatment of Human Umbilical Cord-Derived Mesenchymal Stem Cells With interleukin-6 Abolishes Their Growth-Promoting Effect on Gastric Cancer Cells. In *Int J Mol Med* on 2015 Feb by Mei Wang, Jie Cai et al..PMID:25483835, , (2015)

[PMID:25483835](#)

el at., Aquaporin-4 mediates communication between astrocyte and microglia: Implications of neuroinflammation in experimental Parkinson's disease. In *Neuroscience*. On 2016 Mar 11 by Sun H, Liang R et al..PMID:26774050, , (2016)

[PMID:26774050](#)

el at., Inhibitory effects of eugenol on RANKL-induced osteoclast formation via attenuation of NF- κ B and MAPK pathways. In *Connect Tissue Res* on 2015 Jun by Vishwa Deepak, Abe Kasonga et al..PMID:25405641, , (2015)

[PMID:25405641](#)

el at., Growth inhibition effects of ent-11 δ O⁻-hydroxy-15-oxo-kaur-16-en-19-oic-acid on colorectal carcinoma cells and colon carcinoma-bearing mice. In *Mol Med Rep* on 2016 Apr by Hua Ye, Qiong Wu et al..PMID:2693577, , (2016)

[PMID:2693577](#)

et al., Knockdown of versican V1 induces a severe inflammatory response in LPS-induced acute lung injury via the TLR2-NF- κ B signaling pathway in C57BL/6J mice. In Mol Med Rep on 2016 Jun by Lulu Xu , Tao Xue et al..PMID: 27109786 , , (2016)

[PMID:27109786](#)

et al., Establishment and characterization of bortezomib-resistant U266 cell line: Constitutive activation of NF- κ B-mediated cell signals and/or alterations of ubiquitylation-related genes reduce bortezomib-induced apoptosis. In BMB Rep on 2014 May by Juwon Park, Eun-Kyung Bae et al..PMID: 24286313, , (2014)

[PMID:24286313](#)

et al., Enhanced Survival of Retinal Ganglion Cells Is Mediated by Microglial Cell-Derived PEDF. In Exp Eye Res on 2014 Oct by Jan Darius Unterlauff, Thomas Claudepierre et al..PMID: 25128578, , (2014)

[PMID:25128578](#)

et al., Spautin-1 ameliorates acute pancreatitis via inhibiting impaired autophagy and alleviating calcium overload. In Mol Med on 2016 Oct by Juan Xiao , Xueping Feng et al..PMID:27579473, , (2016)

[PMID:27579473](#)

et al., Fluoxetine protects against IL-1 β -induced neuronal apoptosis via downregulation of p53. In Neuropharmacology on 2016 Aug et al..PMID:Han Shan, Yaqi Bian et al..PMID:26976669, , (2016)

[PMID:26976669](#)

et al., The Effects of Xiangqing Anodyne Spray on treating acute soft-Tissue injury mainly depend on suppressing activations of AKT and p38 pathways. In Evid Based Complement Alternat Med on 2016 by Shudong Wang , Tao Li et al..PMID:27190541, , (2016)

[PMID:27190541](#)

et al., Protodioscin ameliorates fructose-induced renal injury via inhibition of the mitogen activated protein kinase pathway. In Phytomedicine on 2016 Nov 15 by Jinyang Shen, Xiaolin Yang et al..PMID:27765371, , (2016)

[PMID:27765371](#)

et al., Enhancement of anti-acne effect of Scutellaria baicalensis extract by fermentation with symbiotic fungus Penicillium decumbens. In J Biosci Bioeng

on 2020 Nov by Xiaojing Zhu, Yue Mao, et al..PMID: 32747300, , (2020)

[PMID:32747300](#)

et al., Costunolide improved dextran sulfate sodium-induced acute ulcerative colitis in mice through NF- κ B, STAT1/3, and Akt signaling pathways. In Int Immunopharmacol on 2020 Jul by Fan Xie, Hai Zhang, et al..PMID:32413737, , (2020)

[PMID:32413737](#)

et al., The CREB/KMT5A complex regulates PTP1B to modulate high glucose-induced endothelial inflammatory factor levels in diabetic nephropathy. In Cell Death Dis

on 2021 Mar 29 by Ting Huang, Xue Li,et al..PMID:33782381, , (2021)

[PMID:33782381](#)

et al., Vanillin Attenuates Proinflammatory Factors in a tMCAO Mouse Model via Inhibition of TLR4/NF- κ B Signaling Pathway. In Neuroscience on 2022 May 21 by Ping Wang, Chunyi Li, et al..PMID: 35276304, , (2022)

[PMID:35276304](#)

et al., Vitexin Protects against Dextran Sodium Sulfate-Induced Colitis in Mice and Its Potential Mechanisms. In J Agric Food Chem on 2022 Sep 28 by Jing Zhang, Feilin Liang,et al..PMID: 36124900, , (2022)

[PMID:36124900](#)

et al., The A-to-I editing of KPC1 promotes intrahepatic cholangiocarcinoma by attenuating proteasomal processing of NF- κ B p105 to p50. In J Exp Clin Cancer Res

on 2022 Dec 8 by Chengming Gao, Guangming Zhou, et al..PMID:36476255, , (2022)

[PMID:36476255](#)

el at., Targeting HMGB1-NFkb Axis and miR-21 by Glycyrrhizin: Role in Amelioration of Corneal Injury in a Mouse Model of Alkali Burn. In Front Pharmacol on 2022 May 2 by Peihong Wang, Peng Hao, et al..PMID:35586052, , (2022)

[PMID:35586052](#)

el at., Pharmacological inhibition of protein S-palmitoylation suppresses osteoclastogenesis and ameliorates ovariectomy-induced bone lossInJ Orthop TranslatOn2023 Jul 17byLinghui Ma?1?2,?Liwei Zhang et al..PMID: 37521493, , (2023)

[PMID:37521493](#)

el at., The BCL-2 inhibitor APG-2575 resets tumor-associated macrophages toward the M1 phenotype, promoting a favorable response to anti-PD-1 therapy via NLRP3 activationInCell Mol ImmunolOn2024 JanbyFan Luo?#?1,?Han Li et al..PMID:?38062129, , (2023)

[PMID:38062129](#)

el at., Cytoplasmic Escape of Mitochondrial DNA Mediated by Mfn2 Downregulation Promotes Microglial Activation via cGas-Sting Axis in Spinal Cord Injury. In Adv Sci (Weinh) on 2024 Jan by Fei-Long Wei, Tian-Fu Wang,et al..PMID:38009491, , (2024)

[PMID:38009491](#)

Xiuli Zhang;Hao Zhang;Jingting Wang;Yangyi Chen;Jiumao Lin;Qingshui Wang;Cheng Wu;Hui Chen;Yao Lin el at., Curcumin attenuates ulcerative colitis via regulation of Sphingosine kinases 1/NF-kB signaling pathway., , (2025)

[PMID:39832759](#)

Qu Chen;Lingyan Ma;Yang Wen;Wentao Lyu;Minjie Yu;Hua Yang;Yingping Xiao el at., The Effect of Clostridium butyricum-Derived Lipoteichoic Acid on Lipopolysaccharide-Stimulated Porcine Intestinal Epithelial Cells., , (2025)

[PMID:39749788](#)

el at., Differential apoptosis gene expressions of rhabdomyosarcoma cells in response to enterovirus 71 infection.In BMC Infect Dis on 2012 Nov 28 by Shi W, Li X,et al..PMID:23191987, , (2012)

[PMID:23191987](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.