EGFR(Phospho-Ser1070) Antibody

Catalog No: #11080

Package Size: #11080-1 50ul #11080-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$		4.0	
	escri	ntı	nn
$\boldsymbol{\nu}$	COUL	μu	ווט


Product Name	EGFR(Phospho-Ser1070) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of EGFR only when phosphorylated at serine1070.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 1070 (R-Y-S(p)-S-D) derived from Human EGFR.
Target Name	EGFR
Modification	Phospho
Other Names	ERBB1; Receptor protein-tyrosine kinase ErbB-1; kinase EGFR
Accession No.	Swiss-Prot: P00533NCBI Protein: NP_005219.2
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 175kd

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from HUVEC cells untreated or treated with EGF using EGFR(Phospho-Ser1070) Antibody #11080.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using EGFR(Phospho-Ser1070) Antibody #11080(left) or the same antibody preincubated with blocking peptide(right).

Background

Receptor for EGF, but also for other members of the EGF family, as TGF-a, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and increases the interaction of MUC1 with SRC and CTNNB1/beta-catenin.

Feinmesser RL,et al. (1999) J Biol Chem; 274(23): 16168-73.

Gamou S,et al. (1995) FEBS Lett; 357(2): 161-4.

Gamou S, et al. (1994) J Cell Physiol; 158(1): 151-9.

Heisermann GJ, et al. (1988) J Biol Chem; 263(26): 13152-8.

Published Papers

el at., Aldosterone induces myofibroblast EGF secretion to regulate epithelial colonic permeability. In Am J Physiol Cell Physiol on 2013 May 1 by Lluθ sa Mirθ ε, Anna Pθ rez-Bosque, et al..PMID:23467299 , , (2013)

PMID:23467299

Note: This product is for in vitro research use only and is not intended for use in humans or animals.