Rb(Phospho-Ser795) Antibody

Catalog No: #11130

Package Size: #11130-1 50ul #11130-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Rabbit Clonality Polyclonal Isotype IgG Purification Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		
Clonality Polyclonal Isotype IgG Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Product Name	Rb(Phospho-Ser795) Antibody
IgG Purification Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Host Species	Rabbit
Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Clonality	Polyclonal
Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Isotype	IgG
specific antibodies were removed by chromatogramphy using non-phosphopeptide. Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
Applications WB IHC IF Species Reactivity Human Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
Species Reactivity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Specificity The antibody detects endogenous level of Rb only when phosphorylated at serine 795. Immunogen Type Peptide-KLH Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Applications	WB IHC IF
Immunogen Type Peptide-KLH Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Conjugates Unconjugated Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Species Reactivity	Human
Immunogen Description Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb. Unconjugated Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Specificity	The antibody detects endogenous level of Rb only when phosphorylated at serine 795.
Conjugates Unconjugated Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Immunogen Type	Peptide-KLH
Target Name Rb Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Immunogen Description	Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb.
Modification Phospho Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Conjugates	Unconjugated
Other Names P105-RB; PP105; PP110; RB-1; RB1 Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Target Name	Rb
Accession No. Swiss-Prot: P06400NCBI Protein: NP_000312.2 Concentration 1.0mg/ml Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Modification	Phospho
Concentration 1.0mg/ml Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Other Names	P105-RB; PP105; PP110; RB-1; RB1
Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Accession No.	Swiss-Prot: P06400NCBI Protein: NP_000312.2
sodium azide and 50% glycerol.	Concentration	1.0mg/ml
	Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
Storage Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.		sodium azide and 50% glycerol.
	Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 110kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from K562 cells untreated(lane 1) or treated with serum(lane 2) using Rb(Phospho-Ser795) Antibody #11130.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Rb(Phospho-Ser795) Antibody #11130(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using Rb(Phospho-Ser795) Antibody #11130.

Background

Key regulator of entry into cell division that acts as a tumor suppressor. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, SUV420H1 and SUV420H2, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity. Fang Sun , Hanjiang Fu , Qin Liu, et al. (2008) Downregulation of CCND1 and CDK6 by miR-34a induces cell cyclearrest. FEBS Letters, 582:1564

Published Papers

Fang Sun, Hanjiang Fu, Qin Liu el at., Downregulation of CCND1 and CDK6 by miR-34a induces cell cyclearrest., FEBS Letters, 582:1564η— C1568. (2008)

PMID:18406353

Heidi Braumuller, ThomasWieder, Ellen Brenner el at., T-helper-1-cell cytokines drive cancer into senescence., NATURE, 494:361-366(2013)

PMID:23376950

el at., Downregulation of CCND1 and CDK6 by miR-34a Induces Cell Cycle Arrest.In FEBS Lett on 2008 Apr 30 by Fang Sun, Hanjiang Fu,et

al..PMID:18406353, , (2008)

PMID:18406353

el at., T-helper-1-cell Cytokines Drive Cancer Into Senescence. In Nature on 2013 Feb 21 by Heidi Braumn Iler, Thomas Wieder, et al..PMID: 23376950, , (2013)

PMID:23376950

el at., EEN regulates the proliferation and survival of multiple myeloma cells by potentiating IGF-1 secretion.In Biochem Biophys Res Commun on 2014 May 2 by Er-Wen Huang, Sheng-Jiang Xue et al..PMID: 24704450,, (2014)

PMID:24704450

el at., Nuclear distribution of claudin-2 increases cell proliferation in human lung adenocarcinoma cells.In Biochim Biophys Acta on 2014 Sep by Akira Ikari, Ryo Watanabe et al..PMID: 24907662, , (2014)

PMID:24907662

Note: This product is for in vitro research use only and is not intended for use in humans or animals.