# cdc25A(Phospho-Ser76) Antibody

Catalog No: #11138

Package Size: #11138-1 50ul #11138-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| D | esc | rip | otio | on |
|---|-----|-----|------|----|
|   |     |     |      |    |

| Product Name          | cdc25A(Phospho-Ser76) Antibody   |
|-----------------------|--|
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.       |
|                       | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
|                       | specific antibodies were removed by chromatogramphy using non-phosphopeptide.                          |
| Applications          | WB IHC   |
| Species Reactivity    | Human;Mouse;Rat  |
| Specificity           | The antibody detects endogenous level of cdc25A only when phosphorylated at serine 76.                 |
| Immunogen Type        | Peptide-KLH  |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 76 (M-G-S(p)-S-E) derived from Human cdc25A.    |
| Conjugates            | Unconjugated   |
| Target Name           | cdc25A   |
| Modification          | Phospho  |
| Other Names           | CDC25M3; MPIP1;  |
| Accession No.         | Swiss-Prot: P30304NCBI Protein: NP_001780.2  |
| Concentration         | 1.0mg/ml   |
| Formulation           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%   |
|                       | sodium azide and 50% glycerol.   |
| Storage               | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.              |
|                       |  |


# **Application Details**

Predicted MW: 70kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

### **Images**



Western blot analysis of extracts from Hela cells untreated or treated with TNF-a and CA using cdc25A(Phospho-Ser76) Antibody #11138.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using cdc25A(Phospho-Ser76) Antibody #11138(left) or the same antibody preincubated with blocking peptide(right).

# Background

Tyrosine protein phosphatase which functions as a dosage-dependent inducer of mitotic progression. Directly dephosphorylates CDC2 and stimulates its kinase activity. Also dephosphorylates CDK2 in complex with cyclin E, in vitro.

Jin J, et al. (2003) Genes 17(24): 3062-74.

Shimuta K, et al. (2002) EMBO J 21(14): 3694-703.

#### **Published Papers**

Shunfu Piao, Su-Jin Lee, Yongbin Xu el at., CK1ε" θ targets Cdc25A for ubiquitin-mediated proteolysis under normal conditions and in response to checkpoint activation, Cell Cycle, 10(3):531-537(2011)

#### PMID:21252624

el at., CK10 targets Cdc25A for ubiquitin-mediated proteolysis under normal conditions and in response to checkpoint activation. In Cell Cycle on 2011 Feb 1 by Shunfu Piao, Su-Jin Lee, et al..PMID: 21252624, (2011)

PMID:21252624

Note: This product is for in vitro research use only and is not intended for use in humans or animals.