PKR(Phospho-Thr446) Antibody

Catalog No: #11280

Package Size: #11280-1 50ul #11280-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$		4.0	
	escri	nti	าท
$\boldsymbol{\nu}$	COUL	Pur	ווע


Product Name	PKR(Phospho-Thr446) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.	
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho	
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.	
Applications	WB	
Species Reactivity	Hu	
Specificity	The antibody detects endogenous level of PKR only when phosphorylated at threonine 446.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around phosphorylation site of threonine 446 (K-R-T(p)-R-S) derived from Human PKR.	
Target Name	PKR	
Modification	Phospho	
Other Names	ADRB2; E2AK2; EIF2AK2; EIF2aK; PRKR	
Accession No.	Swiss-Prot: P19525 NCBI Protein: NP_001129123.1	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

Application Details

Predicted MW: 68kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from K562 cells using PKR(Phospho-Thr446) Antibody #11280(Lane 2) and the same antibody preincubated with blocking peptide(Lane1).

Background

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection.

Abujiang Pataer, et,al. (2002) Cancer Res; 62: 2239.

K. D. Ryman, et,al. (2005) J. Virol; 79: 1487 - 1499.

Susana Guerra, et,al. (2006) J. Biol. Chem; 281: 18734 - 18745.

Published Papers

Suzette Laing, Guohui Wang, Tamara Briazova el at., Airborne Particulate Matter Selectively Activates Endoplasmic Reticulum Stress Response in the Lung and Liver Tissues., Amerian Journal of Physiol Cell Physiol, 299(4):C736-749. doi:10.1152/ajpcell.00529.2009(2010)

PMID:20554909

el at., Airborne particulate matter selectively activates endoplasmic reticulum stress response in the lung and liver tissues. In Am J Physiol Cell Physiol on 2010 Oct by Suzette Laing, Guohui Wang, et al..PMID: 20554909

, , (2010)

PMID:20554909

el at., Fluoxetine Inhibits NLRP3 Inflammasome Activation: Implication in Depression.In Int J Neuropsychopharmacol.On 2016 Sep 21 by Du RH, Tan J et al..PMID:27207922, , (2016)

PMID:27207922

Note: This product is for in vitro research use only and is not intended for use in humans or animals.