MEK1(Phospho-Thr291) Antibody

Catalog No: #11294

Package Size: #11294-1 50ul #11294-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| _ | | | | |
|---|-----|-------|-----|-----|
| | esc | ~rii | nti | nη |
| - | -0 | 71 II | ווע | וטי |
| | | | | |


| Product Name | MEK1(Phospho-Thr291) Antibody | |
|-----------------------|--|--|
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Isotype | IgG | |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. | |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho | |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. | |
| Applications | WB | |
| Species Reactivity | Human;Mouse;Rat | |
| Specificity | The antibody detects endogenous level of MEK1 only when phosphorylated at threonine 291. | |
| Immunogen Type | Peptide-KLH | |
| Immunogen Description | Peptide sequence around phosphorylation site of threonine 291 (P-R-T(p)-P-G derived from Human MEK1. | |
| Conjugates | Unconjugated | |
| Target Name | MEK1 | |
| Modification | Phospho | |
| Other Names | ERK activator kinase 1; MAP kinase kinase 1; MAP2K1; MAPK/ERK kinase 1; MAPKK 1 | |
| Accession No. | Swiss-Prot: Q02750NCBI Protein: NP_002746.1 | |
| Concentration | 1.0mg/ml | |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% | |
| | sodium azide and 50% glycerol. | |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. | |
| | | |

Application Details

Predicted MW: 45kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from HepG2 cells untreated or treated with PMA using MEK1(Phospho-Thr291) Antibody #11294.

Backgroun<u>d</u>

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates ERK1 and ERK2 MAP kinases.

Kevin D. Burroughs, et,al. (2003) Mol. Cancer Res; 1: 312.

Michael J. Piatelli, et,al. (2002) J. Biol. Chem; 277: 12144 - 12150.

Margaret M. Morgan, et,al. (2001) J. Immunol; 167: 5708.

Herbert Schramek, et,al. (2003) Am J Physiol Cell Physiol, ; 285: C652 - C661.

Published Papers

Hanqian Xu, Gan Zhao, Xiaoxi Huang el at., CD40-expressing plasmid induces anti-CD40 antibody and enhances immune responses to DNA vaccination., The Journal of Gene Medicine., 12(1)97-106(2010)

PMID:19950201

Akira Ikari, Kosuke Atomi, Keishi Kinjo el at., Magnesium deprivation inhibits a MEK CERK cascade and cell proliferation in renal epithelial Madin-Darby canine kidney cells., Life Sciences, 86: 766n— C773(2010)

PMID:20338184

Note: This product is for in vitro research use only and is not intended for use in humans or animals.