

P38 MAPK(Phospho-Thr180/Tyr182) antibody

Catalog No: #11581



Package Size: #11581-1 50ul #11581-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	P38 MAPK(Phospho-Thr180/Tyr182) antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB;IHC;IF
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of P38 MAPK Protein only when phosphorylated at threonine 180/tyrosine 182.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 180 or tyrosine 182 (E-M-T(p)-G-Y(p)-V-A) derived from Human P38 MAPK.
Conjugates	Unconjugated
Target Name	P38 MAPK
Modification	Phospho
Other Names	RK, p38, CSBP, EXIP, Mxi2
Concentration	1.0mg/ml

Application Details

Predicted MW: 43kd


Western blotting: 1: 500-1: 1000 IHC: 1:100-500 IF: 1:100-500

Images




Western blot analysis of extracts from HeLa cells untreated or treated with TNF α + CA using P38 MAPK(Phospho-Thr180/Tyr182) Antibody #11581.


Western blotting analysis using P38
MAPK(Phospho-Thr180/Tyr182) antibody #11581.



Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. P38 (phospho Thr180/Y182) Polyclonal Antibody was diluted at 1:200.



Immunofluorescence analysis of Hela cell. P38 (phospho Thr180/Y182) Polyclonal Antibody(green) was diluted at 1:200.



Background

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK14 is one of the four p38 MAPKs which play an important role in the cascades of cellular responses evoked by extracellular stimuli such as proinflammatory cytokines or physical stress leading to direct activation of transcription factors. Accordingly, p38 MAPKs phosphorylate a broad range of proteins and it has been estimated that they may have approximately 200 to 300 substrates each. Some of the targets are downstream kinases which are activated through phosphorylation and further phosphorylate additional targets. RPS6KA5/MSK1 and RPS6KA4/MSK2 can directly phosphorylate and activate transcription factors such as CREB1, ATF1, the NF-kappa-B isoform RELA/NFKB3, STAT1 and STAT3, but can also phosphorylate histone H3 and the nucleosomal protein HMGN1. RPS6KA5/MSK1 and RPS6KA4/MSK2 play important roles in the rapid induction of immediate-early genes in response to stress or mitogenic stimuli, either by inducing chromatin remodeling or by recruiting the transcription machinery. On the other hand, two other kinase targets, MAPKAPK2/MK2 and MAPKAPK3/MK3, participate in the control of gene expression mostly at the post-transcriptional level, by phosphorylating ZFP36 (tristetraprolin) and ELAVL1, and by regulating EEF2K, which is important for the elongation of mRNA during translation. MKNK1/MNK1 and MKNK2/MNK2, two other kinases activated by p38 MAPKs, regulate protein synthesis by phosphorylating the initiation factor EIF4E2. MAPK14 interacts also with casein kinase II, leading to its activation through autophosphorylation and further phosphorylation of TP53/p53. In the cytoplasm, the p38 MAPK pathway is an important regulator of protein turnover. For example, CFLAR is an inhibitor of TNF-induced apoptosis whose proteasome-mediated degradation is regulated by p38 MAPK phosphorylation. In a similar way, MAPK14 phosphorylates the ubiquitin ligase SIAH2, regulating its activity towards EGLN3. MAPK14 may also inhibit the lysosomal degradation pathway of autophagy by interfering with the intracellular trafficking of the transmembrane protein ATG9. Another function of MAPK14 is to regulate the endocytosis of membrane receptors by different mechanisms that impinge on the small GTPase RAB5A. In addition, clathrin-mediated EGFR internalization induced by inflammatory cytokines and UV irradiation depends on MAPK14-mediated phosphorylation of EGFR itself as well as of RAB5A effectors. Ectodomain shedding of transmembrane

proteins is regulated by p38 MAPKs as well. In response to inflammatory stimuli, p38 MAPKs phosphorylate the membrane-associated metalloprotease ADAM17. Such phosphorylation is required for ADAM17-mediated ectodomain shedding of TGF- α family ligands, which results in the activation of EGFR signaling and cell proliferation. Another p38 MAPK substrate is FGFR1. FGFR1 can be translocated from the extracellular space into the cytosol and nucleus of target cells, and regulates processes such as rRNA synthesis and cell growth. FGFR1 translocation requires p38 MAPK activation. In the nucleus, many transcription factors are phosphorylated and activated by p38 MAPKs in response to different stimuli. Classical examples include ATF1, ATF2, ATF6, ELK1, PTPRH, DDIT3, TP53/p53 and MEF2C and MEF2A. The p38 MAPKs are emerging as important modulators of gene expression by regulating chromatin modifiers and remodelers. The promoters of several genes involved in the inflammatory response, such as IL6, IL8 and IL12B, display a p38 MAPK-dependent enrichment of histone H3 phosphorylation on 'Ser-10' (H3S10ph) in LPS-stimulated myeloid cells. This phosphorylation enhances the accessibility of the cryptic NF- κ B-binding sites marking promoters for increased NF- κ B recruitment. Phosphorylates CDC25B and CDC25C which is required for binding to 14-3-3 proteins and leads to initiation of a G2 delay after ultraviolet radiation. Phosphorylates TIAR following DNA damage, releasing TIAR from GADD45A mRNA and preventing mRNA degradation. The p38 MAPKs may also have kinase-independent roles, which are thought to be due to the binding to targets in the absence of phosphorylation. Protein O-Glc-N-acylation catalyzed by the OGT is regulated by MAPK14, and, although OGT does not seem to be phosphorylated by MAPK14, their interaction increases upon MAPK14 activation induced by glucose deprivation. This interaction may regulate OGT activity by recruiting it to specific targets such as neurofilament H, stimulating its O-Glc-N-acylation. Required in mid-fetal development for the growth of embryo-derived blood vessels in the labyrinth layer of the placenta. Also plays an essential role in developmental and stress-induced erythropoiesis, through regulation of EPO gene expression. Isoform MXI2 activation is stimulated by mitogens and oxidative stress and only poorly phosphorylates ELK1 and ATF2. Isoform EXIP may play a role in the early onset of apoptosis.

Published Papers

Jia-Ping Ruan, M.D., Hong-Xing Zhang et al., EphrinBs/EphBs Signaling Is Involved in Modulation of Spinal Nociceptive Processing through a Mitogen-activated Protein Kinases-dependent Mechanism., *Anesthesiology*, 112:1234-1249(2010)

[PMID:20395829](#)

et al., Astragalus polysaccharides exerts anti-infective activity by inducing human cathelicidin antimicrobial peptide LL-37 in respiratory epithelial cells. *In Phytother Res.* On 2018 Aug by Zhao L, Tan S et al..PMID: 29672953, , (2018)

[PMID:29672953](#)

et al., Molecular mechanisms of somatostatin-mediated intestinal epithelial barrier function restoration by upregulating claudin-4 in mice with DSS-induced colitis. *In Am J Physiol Cell Physiol.* On 2018 Oct 1 by Cai L, Li X et al..PMID: 30020824, , (2018)

[PMID:30020824](#)

et al., Ajudecumin A from *Ajuga ovalifolia* var. *calantha* exhibits anti-inflammatory activity in lipopolysaccharide-activated RAW264.7 murine macrophages and animal models of acute inflammation. *In Pharm Biol.* On 2018 Dec by Zhang H, Ren QC et al..PMID: 31070535, , (2018)

[PMID:31070535](#)

et al., miR-141-5p regulate ATF2 via effecting MAPK1/ERK2 signalling to promote preeclampsia. *Biomed Pharmacother.* In *Biomed Pharmacother* on 2019 May 7 by Wang Y1, Cheng K, et al..PMID: 31075732, , (2019)

[PMID:31075732](#)

et al., HMGN2 regulates non-tuberculous mycobacteria survival via modulation of M1 macrophage polarization. *In J Cell Mol Med* on 2019 Dec by Wang X, Chen S, et al..PMID:31596045, , (2019)

[PMID:31596045](#)

et al., The protective effects of β -sitosterol and vermicularin from *Thamnia vermicularis* (Sw.) Ach. against skin aging in vitro. *In An Acad Bras Cienc* on 2019 Dec 2 by Haiyuan YU, Shen X, et al..PMID:31800700, , (2019)

[PMID:31800700](#)

et al., Attenuating Phosphorylation of p38 MAPK in the Activated Microglia: A New Mechanism for Intrathecal Lidocaine Reversing Tactile Allodynia Following Chronic Constriction Injury in Rats. *In Neurosci Lett* on 2008 Jan 31 by Yi-Wen Gu, Dian-San Su, et al..PMID:18191894, , (2008)

[PMID:18191894](#)

et al., Lidocaine Attenuates Proinflammatory Cytokine Production Induced by Extracellular Adenosine Triphosphate in Cultured Rat Microglia. *In Anesth Analg* on 2010 Sep by Diansan Su, Yang Gu, et al..PMID:20686009, , (2010)

[PMID:20686009](#)

el at., The Anti-Inflammatory Mechanism of Heme oxygenase-1 Induced by Hemin in Primary Rat Alveolar Macrophages. In *Inflammation* on 2012 Jun by Chen Hualin, Xu Wenli, et al..PMID:22160840, , (2012)

[PMID:22160840](#)

el at., Toosendanin Induces Apoptosis Through Suppression of JNK Signaling Pathway in HL-60 Cells. In *Toxicol In Vitro* on 2013 Feb by Jianming Ju, Zhichao Qi, et al..PMID:23111283, , (2013)

[PMID:23111283](#)

el at., Cardamonin Protects Septic Mice From Acute Lung Injury by Preventing Endothelial Barrier Dysfunction. In *J Biochem Mol Toxicol* on 2012 Jul by Zhifeng Wei, Jian Yang, et al..PMID:22696397, , (2012)

[PMID:22696397](#)

el at., Disruption of xCT inhibits cancer cell metastasis via the caveolin-1/ε⁺Y-catenin pathway. In *Oncogene* on 2009 Jan 29 by R-S Chen, Y-M Song, et al..PMID: 19015640, , (2009)

[PMID:19015640](#)

el at., New roles of the fission yeast eIF2^o kinases Hri1 and Gcn2 in response to nutritional stress. In *J Cell Sci* on 2013 Jul 15 by Ruth Marti n, Juan Josθ Berlanga, et al..PMID: 23687372, , (2013)

[PMID:23687372](#)

el at., EphrinBs/EphBs signaling is involved in modulation of spinal nociceptive processing through a mitogen-activated protein kinases-dependent mechanism. In *Anesthesiology*

on 2010 May by Jia-Ping Ruan, Hong-Xing Zhang, et al..PMID: 20395829, , (2010)

[PMID:20395829](#)

el at., Astragaloside IV Inhibits Isoprenaline- ζ ½-induced Cardiac Fibrosis by Targeting the Reactive Oxygen Species/Mitogeni ζ ³ctivated Protein Kinase Signaling Axis. In *Mol Med Rep* on 2017 Apr by Hongliang Dai ,Guizhi Jia ,et al..PMID:28260010, , (2017)

[PMID:28260010](#)

el at., Ellagic Acid Inhibits RANKL-induced Osteoclast Differentiation by Suppressing the p38 MAP Kinase Pathway. In *Arch Pharm Res* on 2017 Jan by Mpho Rantlha , Travers Sagar, et al..PMID: 27384064, , (2017)

[PMID:27384064](#)

el at., Down-regulation of RPS9 inhibits osteosarcoma cell growth through inactivation of MAPK signaling pathway. In *J Cancer* on 2017 Aug 22 by Dong-Dong Cheng, Bin Zhu, et al..PMID: 28928861, , (2017)

[PMID:28928861](#)

el at., Altered Timing of Riboflavin and Ultraviolet Light Pathogen Inactivation Improves Platelet in Vitro Quality. In *Transfusion* on 2017 Aug by Peter Schubert, Brankica Culibrk, et al..PMID: 28500654, , (2017)

[PMID:28500654](#)

el at., Hepatic SATB1 induces paracrine activation of hepatic stellate cells and is upregulated by HBx. In *Sci Rep*. On 2016 Nov 24 by Gong J, Tu W et al..PMID:27883059, , (2016)

[PMID:27883059](#)

el at., Quercetin regulates the sestrin 2-AMPK-p38 MAPK signaling pathway and induces apoptosis by increasing the generation of intracellular ROS in a p53-independent manner. In *Int J Mol Med* on 2014 Apr by Guen Tae Kim, Se Hee Lee et al..PMID:24535669, , (2014)

[PMID:24535669](#)

el at., Potassium supplementation inhibits IL-17A production induced by salt loading in human T lymphocytes via p38/MAPK-SGK1 pathway. In *Exp Mol Pathol*. On 2016 Jun by Wen W, Wan Z et al..PMID:27020669, , (2016)

[PMID:27020669](#)

el at., The activation of p38 MAPK limits the abnormal proliferation of vascular smooth muscle cells induced by high sodium concentrations. In *Int J Mol Med* On 2016 Jan by Yan Wu , Juan Zhou et al..PMID:26530729, , (2016)

[PMID:26530729](#)

el at., Regulation of T lymphocyte activation by microRNA-21.In Mol Immunol on 2014 Jun by Lu Wang , Liangqiang He et al..PMID: 24631982, , (2014)

[PMID:24631982](#)

el at., Piperine alleviates osteoclast formation through the p38/ci ζ os/NFATc1 signaling axis.In Biofactors on Nov-Dec 2015 by Vishwa Deepak , Marlena C Kruger et al..PMID: 26627060, , (2015)

[PMID:26627060](#)

el at., The Effects of Xiangqing Anodyne Spray on treating acute soft-Tissue injury mainly depend on suppressing activations of AKT and p38 pathways.In Evid Based Complement Alternat Med on 2016 by Shudong Wang , Tao Li et al..PMID:27190541, , (2016)

[PMID:27190541](#)

el at., Insights in dynamic kinome reprogramming as a consequence of MEK inhibition in MLL-rearranged AML.In Leukemia on 2014 Mar by K R Kampen, A Ter Elst et al..PMID:24240200, , (2014)

[PMID:24240200](#)

el at., Cannabinoid receptor agonist WIN55,212-2 and fatty acid amide hydrolase inhibitor URB597 suppress chronic cerebral hypoperfusion-induced neuronal apoptosis by inhibiting c-Jun N-terminal kinase signaling.In Neuroscience 2015 Aug 20 by S-H Su , Y-F Wu et al..PMID:25795598, , (2015)

[PMID:25795598](#)

el at., PiHOG1, a stress regulator MAP kinase from the root endophyte fungus Piriformospora indica, confers salinity stress tolerance in rice plants. In Sci Rep on 2016 Nov 16 by Abhimanyu Jogawat, Jyothilakshmi Vadassery, et al..PMID:27849025, , (2016)

[PMID:27849025](#)

el at., Enhancement of anti-acne effect of Scutellaria baicalensis extract by fermentation with symbiotic fungus Penicillium decumbens. In J Biosci Bioeng on 2020 Nov by Xiaojing Zhu, Yue Mao, et al..PMID: 32747300, , (2020)

[PMID:32747300](#)

el at., Role of 5-HT degradation in acute liver injury induced by carbon tetrachloride. In Eur J Pharmacol on 2021 Oct 5 by Yu-Xin Zhang, Chen Li,et al..PMID:34280394, , (2021)

[PMID:34280394](#)

el at., CD155 Cooperates with PD-1/PD-L1 to Promote Proliferation of Esophageal Squamous Cancer Cells via PI3K/Akt and MAPK Signaling Pathways. In Cancers (Basel) on 2022 Nov 15 by Xiyang Tan, Jie Yang, et al..PMID:36428703, , (2022)

[PMID:36428703](#)

el at., Brevilin A Ameliorates Acute Lung Injury and Inflammation Through Inhibition of NF-κB Signaling via Targeting IKKα/β. In Front Pharmacol on 2022 Jun 14 by Lu Liu, Xian Chen,et al..PMID:35774606, , (2022)

[PMID:35774606](#)

el at., Corilagin ameliorates macrophages inflammation in atherosclerosis through TLR4-NFκB/MAPK pathwayIn HeliyonOn2023 Jun 13byDa Meng , Xin Deng et al..PMID:37383215, , (2023)

[PMID:37383215](#)

el at., Gingerenone A Attenuates Ulcerative Colitis via Targeting IL-17RA to Inhibit Inflammation and Restore Intestinal Barrier Function. In Adv Sci (Weinh) on 2024 Jul by Jian Liang, Weigang Dai,et al..PMID:38639442, , (2024)

[PMID:38639442](#)

Jing Guan;Xin Tong;Yi Zhang;Fan Xu;Yuxin Zhang;Xiurui Liang;Jiaqi Jin;Hongyan Jing;Liuxian Guo;Xinrui Ni;Jihua Fu el at., Nephrotoxicity induced by cisplatin is primarily due to the activation of the 5-hydroxytryptamine degradation system in proximal renal tubules, , (2021)

[PMID:](#)

Yu Cao;Yongju Wen;Zongyuan Zhou;Ruiying Xi;Wen Shuai;Jichao Zhang;Apichart Suksamrarn;Guolin Zhang;Xiao-Xia Lu;Fei Wang el at., Discovery of novel capsaicin analogs as TRPV1 inhibitors for the treatment of idiopathic pulmonary fibrosis., , (2025)

[PMID:39826937](#)

et al., CPU-12, a novel synthesized oxazolo [5, 4-d] pyrimidine derivative, showed superior anti-angiogenic activity. In J Pharmacol Sci on 2015 Sep by Jiping Liu, Ya-Hui Deng et al.. PMID: 26154846, (2015)

[PMID:26154846](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.