

## TrkB(Phospho-Y817) Rabbit mAb

Catalog No: #13384



Package Size: #13384-1 50ul #13384-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

Product Name	TrkB(Phospho-Y817) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	SC0556
Purification	ProA affinity purified
Applications	WB;ICC/IF;IHC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Tyr817 of human TrkB.
Conjugates	Unconjugated
Accession No.	Swiss-Prot#:Q16620
Calculated MW	145 kDa
SDS-PAGE MW	140 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:500-1:2000


ICC/IF: 1:50-1:200

IHC: 1:50-1:200


## Images




All lanes : TrkB(Phospho-Y817) Rabbit mAb at 1/1k dilution  
 Lane 1 : SH-SY5Y whole cell lysates  
 Lane 2 : Mouse brain tissue lysates  
 Lane 3 : Rat brain tissue lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution  
 Predicted band size: 145 kDa  
 Observed band size: 140 kDa  
 Exposure time: 12 seconds




Formalin-fixed;paraffin-embedded human kidney tissue stained for TrkB (Phospho-Y817) using 13384 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed;paraffin-embedded human pancreas tissue stained for TrkB (Phospho-Y817) using 13384 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed;paraffin-embedded mouse pancreas tissue stained for TrkB (Phospho-Y817) using 13384 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence  
TrkB(Phospho-Y817) antibody ICC/IF staining of TrkB(Phospho-Y817) in HeLa cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with antibody at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500. The cells were incubated 30 minutes at room temperature with phalloidin tritc . Nuclei were counterstained with DAPI.

## Background

The Trk proto-oncogene encodes a tyrosine protein kinase, Trk A, also designated Trk gp140, that serves as a receptor for certain neurotrophic factors including nerve growth factor (NGF) and neurotrophin-3 (NT-3). Trk B is a tyrosine kinase gene highly related to Trk A. Trk B expression is confined to tissues within the central and peripheral nervous systems. The brain-derived neurotrophic factor (BDNF) and NT-3, but not NGF, can induce rapid phosphorylation on tyrosine of Trk B gp145, one of the receptors encoded by NTRK2, although BDNF elicits a response at least two orders of magnitude greater than NT-3. Thus it appears that Trk B gp145 may represent a neurotrophic receptor for BDNF and NT-3. The third member of the Trk family of tyrosine kinases, Trk C, encodes a protein designated Trk C gp145 that is preferentially expressed in brain tissue, is equally related to Trk A and Trk B and is a functional receptor for NT-3.

## References

---

1. Nannini M et al. Integrated genomic study of quadruple-WT GIST (KIT/PDGFR $\alpha$ /SDH/RAS pathway wild-type GIST). *BMC Cancer* 14:685 (2014).
2. Zhang HH et al. The BDNF/TrkB signaling pathway is involved in heat hyperalgesia mediated by Cdk5 in rats. *PLoS One* 9:e85536 (2014).

## Published Papers

---

Zhang Qin, Liu Jiayi, Wu Bingfeng, Yuan Ying, Gong Ping, Xiang Lin et al., Lysophosphatidic acid-induced YAP activation regulates osteogenesis of MC3T3-E1, *Tissue & cell*, (2025)

[PMID:40902513](#)

---

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.