

AKT1(Phospho-S124) Rabbit mAb

Catalog No: #13423



Package Size: #13423-1 50ul #13423-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	AKT1(Phospho-S124) Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JJ08-46
Purification	ProA affinity purified
Applications	WB;ICC/IF;IHC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser124 of human AKT1.
Conjugates	Unconjugated
Other Names	AKT 1 antibody AKT antibody AKT1 antibody AKT1_HUMAN antibody MGC99656 antibody PKB antibody PKB-ALPHA antibody PRKBA antibody Protein Kinase B Alpha antibody Protein kinase B antibody Proto-oncogene c-Akt antibody RAC Alpha antibody RAC antibody RAC-alpha serine/threonine-protein kinase antibody RAC-PK-alpha antibody
Accession No.	Swiss-Prot#:P31749
Calculated MW	56 kDa
SDS-PAGE MW	56 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:2000

ICC/IF: 1:50-1:200

IHC: 1:50-1:200

Images



All lanes : AKT1(Phospho-S124) Rabbit mAb at 1/1000 dilution

Lane 1 : MCF7 whole cell lysates treated with λ pp for 1 hour

Lane 2 : MCF7 whole cell lysates

Lane 3 : PC12 whole cell lysates

Lane 4 : NIH/3T3 whole cell lysates

Lysates/proteins at 20 μ g per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution


Predicted band size: 56 kDa

Observed band size: 56 kDa

Exposure time: 12 seconds



Formalin-fixed;paraffin-embedded human uterus cancer tissue stained for AKT1 (Phospho-S124) using 13423 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence AKT1 (Phospho-S124) antibody (13423)
ICC/IF staining of AKT1(Phospho-S124) in NIH/3T3 cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with 13423 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit;used at a dilution of 1/500. Nuclei were counterstained with DAPI.

Background

The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKB β or RacPK- β b) and Akt 3 (also designated PKB γ ; or thymoma viral proto-oncogene 3), which exhibit sequence homology with the protein kinase A and C families and are encoded by the c-Akt proto-oncogene. All members of the Akt family have a pleckstrin homology domain. Akt1 and Akt2 are activated by PDGF stimulation. Activation is dependent on PDGFR- β Tyr residues 740 and 751, which bind the subunit of the phosphatidylinositol 3-kinase (PI 3-kinase) complex. Activation of Akt1 by Insulin or Insulin-growth factor-1(IGF-1) results in phosphorylation of both Thr 308 and Ser 473. Phosphorylation of both residues is important to generate a high level of Akt1 activity. The phosphorylation of Thr 308 is not dependent on phosphorylation of Ser 473 *in vivo*. Thus, Akt proteins become phosphorylated and activated in Insulin/IGF-1-stimulated cells by an upstream kinase(s). The activation of Akt1 and Akt2 is inhibited by the PI kinase inhibitor wortmannin, suggesting that the protein signals downstream of the PI kinases.

References

1. Lv P et al. Treatment with the herbal medicine, naoxintong improves the protective effect of high-density lipoproteins on endothelial function in patients with type 2 diabetes. *Mol Med Rep* 13:2007-16 (2016).
2. Xu L et al. MicroRNA-7-regulated TLR9 signaling-enhanced growth and metastatic potential of human lung cancer cells by altering the phosphoinositide-3-kinase, regulatory subunit 3/Akt pathway. *Mol Biol Cell* 24:42-55 (2013).

Published Papers

Shanxing Dango^o Xiaoxuan Tango^o Qingyun Lio^o June Wu^o Yue Huo^o Xiqing Lio^o Danong Cao^o Chuwen Fengo^o Yehong Suno^o Jianmei Huang^o Wencong Song^o Xuelian Du et al., Xiao-Liu Tang (XLT), a traditional Chinese medicine formula that suppresses the progression of cervical cancer by inducing apoptosis and inhibiting cell migration, *Journal of ethnopharmacology*, (2025)

PMID:41344524

Note: This product is for in vitro research use only and is not intended for use in humans or animals.