Product Datasheet

Smad2(Phospho-S255) Rabbit mAb

Catalog No: #13429

Package Size: #13429-1 50ul #13429-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

- -		1		
пре	\sim rı	OT.	$\boldsymbol{\cap}$	n
Des	ווטי	บแ	v	

•	
Product Name	Smad2(Phospho-S255) Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JF0882
Purification	ProA affinity purified
Applications	WB,IHC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser255 of human Smad2.
Conjugates	Unconjugated
Other Names	Drosophila, homolog of, MADR2 antibody hMAD-2 antibody HsMAD2 antibody JV18 antibody JV18-1
	antibody JV181 antibody MAD antibody MAD homolog 2 antibody MAD Related Protein 2 antibody
	Mad-related protein 2 antibody MADH2 antibody MADR2 antibody MGC22139 antibody MGC34440
	antibody Mother against DPP homolog 2 antibody Mothers against decapentaplegic homolog 2 antibody
	Mothers against decapentaplegic, Drosophila, homolog of, 2 antibody Mothers against DPP homolog 2
	antibody OTTHUMP00000163489 antibody Sma and Mad related protein 2 antibody Sma- and Mad-related
	protein 2 MAD antibody SMAD 2 antibody SMAD family member 2 antibody SMAD, mothers against DPP
	homolog 2 antibody SMAD2 antibody SMAD2_HUMAN antibody
Accession No.	Swiss-Prot#:Q15796
Calculated MW	52 kDa
SDS-PAGE MW	60 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:2000 IHC: 1:50-1:200

Background

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of TGFβ/BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGFβ and Activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to Activin/TGFβ signaling by interfering with TGFβ-mediated phosphorylation of other Smad proteins.

References

1. Ungefroren H et al. Rac1b negatively regulates TGF-?1-induced cell motility in pancreatic ductal epithelial cells by suppressing Smad signalling.

Oncotarget 5:277-90 (2014).

2. Harazono Y et al. miR-655 Is an EMT-suppressive MicroRNA targeting ZEB1 and TGFBR2. PLoS One 8:e62757 (2013).

Published Papers

el at., Gastrodin attenuates renal injury and collagen deposition via suppression of the TGF- β 1/Smad2/3 signaling pathway based on network pharmacology analysis InFront PharmacolOn2023 Jan 17byYing Wen , Xiuli Zhang et al..PMID:36733505, , (2023)

PMID:36733505

Note: This product is for in vitro research use only and is not intended for use in humans or animals.