Product Datasheet

Myc Antibody

Catalog No: #21034

Package Size: #21034-1 50ul #21034-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$		4.0	
	escri	ntı	nn
$\boldsymbol{\nu}$	COUL	μu	ווט

Product Name	Myc Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB IHC	
Species Reactivity	Human;Mouse;Rat	
Specificity	The antibody detects endogenous level of total Myc protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa.56~60 (L-P-T-P-P) derived from Human Myc.	
Conjugates	Unconjugated	
Target Name	Мус	
Other Names	с-тус	
Accession No.	Swiss-Prot: P01106NCBI Protein: NP_002458.2	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	


Application Details

Predicted MW: 60kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from MCF cells using Myc(Ab-58) Antibody #21034.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Myc(Ab-58) Antibody #21034(left) or the same antibody preincubated with blocking peptide(right).

Background

Myc proto-oncogene encodes nuclear DNA-binding phosphoproteins that are involved in the regulation of gene expression and DNA replication during cell growth and differentiation. Myc encodes a protein of 65 kDa which is expressed in almost all normal and transformed cells. The expression correlates with the proliferation state of the cells. Transcription is repressed in quiescent or terminally differentiated cells. Expression of Myc is generally induced after mitogenic stimulation of cells or serum induction. Myc therefore is an important positive regulator of cell growth and proliferation. Myc has been demonstrated also to be a potent inducer of apoptosis when expressed in the absence of serum or growth factors. Apoptosis may serve also as a protective mechanism to prevent tumorigenicity elicited by deregulated Myc expression. Sequences of the Myc oncogene have been highly conserved throughout evolution, from Drosophila to vertebrates

Jin Z, et al. (2004) J Biol Chem. 279(38): 40209-40219.

Welcker M, et al. (2004) Proc Natl Acad Sci U S A. 101(24): 9085-9090.

Baudino T A, et al. (2001) Mol Cell Biol. 21: 691-702.

Blackwood E M, et al. (1991) Science. 251:1211-1217.

Henriksson M, et al. (1996) Adv Cancer Res. 68: 109-182.

Published Papers

S.Y. Liu, Y.L. Ma, E.H.Y. Lee el at., NMDA receptor signaling mediates the expression of protein inhibitor of activated STAT1 (PIAS1) in rat hippocampus., Neuropharmacology, 65:101-113(2013)

PMID:22982248

el at., NMDA receptor signaling mediates the expression of protein inhibitor of activated STAT1 (PIAS1) in rat hippocampus. In Neuropharmacology on 2013 Feb by S Y Liu, Y L Ma, et al..PMID: 22982248

, , (2013)

PMID:22982248

el at., MiRNA-29a as a tumor suppressor mediates PRIMA-1Met-induced anti-myeloma activity by targeting c-Myc. In Oncotarget on 2016 Feb 9 by Manujendra N Saha, Jahangir Abdi, et al..PMID: 26771839, , (2016)

PMID:26771839

Note: This product is for in vitro research use only and is not intended for use in humans or animals.