

STAT6 Antibody

Catalog No: #21050

Package Size: #21050-1 50ul #21050-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	STAT6 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of total STAT6 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.639~643 (R-G-Y-V-P) derived from Human STAT6.
Conjugates	Unconjugated
Target Name	STAT6
Other Names	IL-4 Stat
Accession No.	Swiss-Prot: P42226NCBI Protein: NP_003144.3
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 110kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from HepG2 cells using STAT6(Ab-641) Antibody #21050.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using STAT6(Ab-641) Antibody #21050(left) or the same antibody preincubated with blocking peptide(right).

Background

STAT6 is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor-associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. By screening an embryonic lung fibroblast cDNA library with a wildtype STAT6 probe, 2 variant cDNAs were identified, which were termed STAT6B and STAT6C, encoding an N-terminal 110-amino acid truncation and a 27-amino acid deletion in the SH2 domain, respectively. All the three variants are ubiquitously expressed with STAT6B expression greatest in spleen and STAT6C expression greatest in lung. STAT6B resembles an attenuated STAT6, but that STAT6C inhibits IL-4-mediated mitogenesis and cell surface antigen expression, and is not tyrosine phosphorylated

Nelms K, et al. (1999) Annu Rev Immunol. 17:701-738.

Malabarba M G, et al. (1996) Biochem. J. 319:865-872.

Hou J, et al. (1994) Science. 265:1701-1706.

Quelle F W, et al. (1995) Mol Cell Biol. 15: 3336-3343.

Takeda K, et al. (1996) Nature. 380: 627-630.

Published Papers

el at., FOXQ1/NDRG1 axis exacerbates hepatocellular carcinoma initiation via enhancing crosstalk between fibroblasts and tumor cells.In Cancer Lett.

On 2018 Mar 28 by Luo Q, Wang CQ et al..PMID: 29248714, , (2018)

[PMID:29248714](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.