

p53 Antibody

Catalog No: #21083

Package Size: #21083-1 50ul #21083-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	p53 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB;IHC;IF;IP;ELISA
Species Reactivity	Human
Specificity	The antibody detects endogenous level of total p53 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.4~8 (P-Q-S-D-P) derived from Human p53.
Conjugates	Unconjugated
Target Name	p53
Other Names	Tumor suppressor p53; Phosphoprotein p53; Antigen NY-CO-13; TP53;
Accession No.	Swiss-Prot: P04637NCBI Protein: NP_000537.3
Calculated MW	44kDa
SDS-PAGE MW	53kDa
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

WB 1:500-1:2000; IHC 1:50-1:200; IF 1:50-1:200; ELISA 1:5000-1:20000; IP 1:10-1:200;

Images



Western blot analysis of extracts from MDA cells using p53(Ab-6) Antibody #21083.

Background

p53 is a nuclear protein which plays an essential role in the regulation of cell cycle specifically in the transition from G0 to G1. It is found in very low levels in normal cells however in a variety of transformed cell lines in high amounts and believed to contribute to transformation and malignancy. The open reading frame of p53 is 393 amino acids long, with the central region (consisting of amino acids from about 100 to 300) containing the DNA-binding domain. This proteolysis-resistant core is flanked by a C-terminal end mediating oligomerization and an N-terminal end containing a strong transcription activation signal. p53 binds as a tetramer to a PBS (p53-Binding Site) and activates the expression of downstream genes that inhibit growth and/or invasion. p53 binds as a tetramer to a p53-binding site (PBS) and to activate the expression of adjacent genes that inhibit growth and/or invasion. Deletion of one or both p53 alleles reduces the expression of tetramers, resulting in decreased expression of the growth inhibitory genes

Lin T, et al. (2005) Nat Cell Biol; 7(2): 165-71.

Vega FM, et al. (2004) Mol Cell Biol; 24(23): 10366-80.

Li J, et al. (2004) J Biol Chem; 279(40): 41275-9.

Wang J, et al. (2004) J Biol Chem; 279(38): 39584-92.

Published Papers

el at., AKT1 ζ• dependent phosphorylation of Niban regulates nucleophosmini ζ• nd MDM2i ζ• ediated p53 stability and cell apoptosis. In EMBO Rep on 2012 Jun 1 by Haitao Ji, Zhiyong Ding, et al..PMID: 22510990

, , (2012)

[PMID:22510990](#)

el at., Down-Regulated Exosomal MicroRNA-221 - 3p Derived From Senescent Mesenchymal Stem Cells Impairs Heart Repair. In Front Cell Dev Biol on 2020 May 5 by Ling Sun, Wenwu Zhu, et al..PMID: 32432109, , (2020)

[PMID:32432109](#)

el at., Long noncoding RNA UCA1 from hypoxia-conditioned hMSC-derived exosomes: a novel molecular target for cardioprotection through miR-873-5p/XIAP axis. In Cell Death Dis on 2020 Aug 10 by Ling Sun, Wenwu Zhu, et al..PMID:32826854, , (2020)

[PMID:32826854](#)

Lei Zhang, Junshan Ruan, Linggeng Yan el at., Xanthatin Induces Cell Cycle Arrest at G2/M Checkpoint and Apoptosis via Disrupting NF-ε" θ 'B Pathway in A549 Non-Small-Cell Lung Cancer Cells., Molecules., 17:3736-3750(2012)

[PMID:22450683](#)

el at., Nordihydroguaiaretic acid inhibits growth of cervical cancer SiHa cells by up-regulating p21.In Oncol Lett on 2011 Jan by Peng Gao, Fei Zhai,et al..PMID:22870140, , (2011)

[PMID:22870140](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.