Tau Antibody

Catalog No: #21093

Package Size: #21093-1 50ul #21093-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


D	es	cri	b.	tic	ıc	1

Product Name	Tau Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were		
	purified by affinity-chromatography using epitope-specific peptide.		
Applications	WB;IHC;IF;ELISA		
Species Reactivity	Human;Mouse;Rat		
Specificity	The antibody detects endogenous level of total Tau protein.		
Immunogen Type	Peptide-KLH		
Immunogen Description	Peptide sequence around aa. 394~398 (Y-K-S-P-V) derived from Human Tau.		
Conjugates	Unconjugated		
Target Name	Tau		
Other Names	Neurofibrillary tangle protein; Paired helical filament-tau;		
Accession No.	Swiss-Prot: P10636NCBI Protein: NP _001116538.1		
Calculated MW	48 62 78 kDa		
SDS-PAGE MW	50-80kDa		
Concentration	1.0mg/ml		
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%		
	sodium azide and 50% glycerol.		
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.		


Application Details

WB 1:500-1:2000; IHC 1:100-1:300; IF 1:50-1:500; ELISA 1:20000;

Images



Western blot analysis of extracts from mouse brain tissue using Tau(Ab-396) Antibody #21093.



Immunohistochemical analysis of paraffin-embedded rat hippocampal region tissue from a model with Alzheimer

Background

Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.

Puig B, et al. (2005) Acta Neuropathol (Berl). 110(3):261-268.

DeGiorgis JA, et al. (2005) J Proteome Res. 4(2): 306-315.

Alonso Adel C, et al. (2004) J Biol Chem. 279(33): 34873-34881.

Published Papers

el at., Tau hyperphosphorylation is associated with memory impairment after exposure to 1.5% isoflurane without temperature maintenance in rats.In Eur J Anaesthesiol on 2010 Sep by Tan W, Cao X, et al..PMID:20485178, (2010)

PMID:20485178

el at., Protective effects of lithium treatment for spatial memory deficits induced by tau hyperphosphorylation in splenectomized rats.In Clin Exp Pharmacol Physiol on 2010 Oct by Tan WF, Cao XZ,et al..PMID:20659131, , (2010)

PMID:20659131

el at., Chronic Cold-Water-Induced Hypothermia Impairs Memory Retrieval and Nepeta Menthoides as a Traditional "Hot" Herb Reverses the Impairment.In Iran J Pharm Res on Winter 2014 by Mohammad Mahdi ,Ahmadian-Attar et al..PMID:24711845, , (2014)

PMID:24711845

el at., Impairment of rat spatial learning and memory in a new model of cold water-induced chronic hypothermia: implication for Alzheimer's disease.In Neurotox Res on 2015 Aug by Mohammad Mahdi Ahmadian-Attari, Leila Dargahi et al..PMID:25782579, , (2015)

PMID:25782579

el at., Capsaicin consumption reduces brain amyloid-beta generation and attenuates Alzheimer's disease-type pathology and cognitive deficits in APP/PS1 mice. In Transl Psychiatry on 2020 Jul 13 by Jun Wang, Bin-Lu Sun, et al.. PMID:32661266, , (2020)

PMID:32661266

el at., Oleocanthal enhances amyloid-ε^ωY clearance from the brains of TgSwDI mice and in vitro across a human blood-brain barrier model. In ACS Chem Neurosci on 2015 Nov 18 by Hisham Qosa, Yazan S Batarseh,et al..PMID:26348065, , (2015)

PMID:26348065

ZhiHao Liu;YuDi Bai;ZhongYuan Yu;HuiYun Li;Jie Liu;ChengRong Tan;GuiHua Zeng;YunFeng Tu;PuYang Sun;YuJuan Jia;JinCai He;YanJiang Wang;XianLe Bu el at., Improving Blood Monocyte Energy Metabolism Enhances Its Ability to Phagocytose Amyloid-β and Prevents Alzheimer's Disease-Type Pathology and Cognitive Deficits, , (2023)

PMID:37316674

el at., Erythropoietin attenuates Alzheimer-like memory impairments and pathological changes induced by amyloid ϵ° Y42 in mice.In Brain Res on

2015 Aug 27 by Yi-Pei Li, Guo-Jun Yang et al..PMID:26049128, , (2015)

PMID:26049128

Meng-Ting Wang;Zi-Cheng Hu;Yang Xiang;Xiao-Qin Zeng;Zhang-Cheng Fei;Jia Chen;Xin-Peng Li;Yu-Peng Zhu;Jun Wang;Yan-Jiang Wang;Zhi-Qiang Xu;Yu-Hui Liu el at., Fingolimod ameliorates amyloid deposition and neurodegeneration in APP/PS1 mouse model of Alzheimer's disease., , (2025)

PMID:40158900

el at., Gypenoside IX restores Akt/GSK-3β pathway and alleviates Alzheimer's disease-like neuropathology and cognitive deficitsInAging (Albany NY)On2023 Dec 12byLing Lei?1?2,?Yong Luo et al..PMID:?38095632, , (2023)

PMID:38095632

Note: This product is for in vitro research use only and is not intended for use in humans or animals.