

Pyk2 Antibody

Catalog No: #21209

Package Size: #21209-1 50ul #21209-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Pyk2 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC IF
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of total Pyk2 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.400~404 (D-I-Y-A-E) derived from Human Pyk2.
Conjugates	Unconjugated
Target Name	Pyk2
Other Names	FADK 2; FAK2; Focal adhesion kinase 2; PTK2B; Proline-rich tyrosine kinase 2 RAFTK
Accession No.	Swiss-Prot: Q14289NCBI Protein: NP_004094.3
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 140kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from 293 cells using Pyk2(Ab-402) Antibody #21209.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Pyk2(Ab-402) Antibody #21209(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using Pyk2(Ab-402) Antibody #21209.

Background

Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation.

Gluck SL, et al. (2004) J Clin Invest; 114(12): 1696-1699

Benzing T, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9784-9789

Tian D, et al. (2002) Mol Cell Biol; 22(8): 2650-2662

Lu Z, et al. (2001) Mol Cell Biol; 21(12): 4016-4031

Krishnan HH, et al. (2006) J Virol; 80(3): 1167-1180

Published Papers

Rui-Fang Liu, Xiao Xu, Jian Huang et al., Down-regulation of miR-517a and miR-517c promotes proliferation of hepatocellular carcinoma cells via targeting Pyk2, Cancer Letters, 329:164-173(2013)

[PMID:23142219](#)

et al., Down-regulation of miR-517a and miR-517c Promotes Proliferation of Hepatocellular Carcinoma Cells via Targeting Pyk2. In Cancer Letters 2013 Feb 28 by Rui-Fang Liu, Xiao Xu, et al.. PMID:23142219, (2013)

[PMID:23142219](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.