Smad2 Antibody

Catalog No: #21323

Package Size: #21323-1 50ul #21323-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

	4.5	
Descri	ntin	n
DUSCH	puo	

Product Name	Smad2 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of total Smad2 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.218~222 (P-E-T-P-P) derived from Human Smad2.
Conjugates	Unconjugated
Target Name	Smad2
Other Names	JV18-1; MADH2; MADR2; Mad-related protein 2; Mothers against DPP homolog 2
Accession No.	Swiss-Prot: Q15796NCBI Protein: NP _001003652.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 60kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from 293 and 3T3 cells using Smad2(Ab-220) Antibody #21323



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Smad2(Ab-220) Antibody #21323(left) or the same antibody preincubated with blocking peptide(right).

Background

Transcriptional modulator activated by TGF-beta and activin type 1 receptor kinase. SMAD2 is a receptor-regulated SMAD (R-SMAD). May act as a tumor suppressor in colorectal carcinoma.

Sang Gyun Kim et al. (2004) Mol Biol Cell. February; 15(2): 420

Published Papers

el at., Distribution of Smad mRNA and proteins in the rat brain.In J Chem Neuroanat. On 2018 Jul by Nakajima T, Hata R et al..PMID: 29196107, , (2018)

PMID:29196107

el at., TGF β1 mediated Smad signaling pathway and EMT in hepatic fibrosis induced by Nano NiO in vivo and in vitro.In Environ Toxicol on 2020 Apr by Zhang Q, Chang X ,et al..PMID: 31737983, , (2020)

PMID:31737983

el at., Temporal and regional patterns of Smad activation in the rat hippocampus following global ischemia.In J Neurol Sci on 2014 Feb 15 by Takayuki Nakajima , Masafumi Yanagihara et al ..PMID:24290497, , (2014)

PMID:24290497

Note: This product is for in vitro research use only and is not intended for use in humans or animals.