Myc Mouse Monoclnal Antibody

Catalog No: #21390

Package Size: #21390-1 50ul #21390-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

<u> </u>	
Descri	ntion
DUSCII	Puon


Product Name	Myc Mouse Monoclnal Antibody
Host Species	Mouse
Clonality	Monoclonal
Clone No.	3.00E+08
Isotype	lgG1
Purification	Antibodies were produced from mice ascites by injecting mice with a monoclonal cell line which was fusioned
	by mouse spleen and SP2/0 myeloma cell. Spleen cells were isolated from mice by immunizing with
	synthetic peptide and KLH conjugates.
Applications	WB
Species Reactivity	Hu
Specificity	This mouse mAb only detects transfected proteins.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence derived from C-terminal (aa. 410-432) of human c-Myc conjugated with KLH.
Target Name	Мус
Other Names	c-myc
Accession No.	Swiss-Prot: P01106NCBI Protein: NP_002458.2
Concentration	1.0mg/ml
Formulation	Supplied in mice ascites.
Storage	Store at 4°C for short term use. Store at -20°C for long term preservation. Avoid freezing and thawing repeatly.

Application Details

Predicted MW: 60kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from 293 cells transfected with recombinant human Myc using Myc mouse mAb #21390.

Background

Myc proto-oncogene encodes nuclear DNA-binding phosphoproteins that are involved in the regulation of gene expression and DNA replication during cell growth and differentiation. Myc encodes a protein of 65 kDa which is expressed in almost all normal and transformed cells. The expression correlates with the proliferation state of the cells. Transcription is repressed in quiescent or terminally differentiated cells. Expression of Myc is generally induced after mitogenic stimulation of cells or serum induction. Myc therefore is an important positive regulator of cell growth and proliferation. Myc has been demonstrated also to be a potent inducer of apoptosis when expressed in the absence of serum or growth factors. Apoptosis may serve also as a protective mechanism to prevent tumorigenicity elicited by deregulated Myc expression. Sequences of the Myc oncogene have been highly conserved throughout evolution, from Drosophila to vertebrates

Baudino T A, et al. (2001) Mol Cell Biol. 21: 691-702.

Blackwood E M, et al. (1991) Science. 251:1211-1217.

Henriksson M, et al. (1996) Adv Cancer Res. 68: 109-182.

Grandori C, et al. (2000) Annu Rev Cell Dev Biol. 16: 653-699.

Published Papers

el at., ERK1/2-dependent phosphorylation and nuclear translocation of PKM2 promotes the Warburg effect. In Nat Cell Biol on 2012 Dec by Weiwei Yang, Yanhua Zheng, et al.. PMID:23178880, , (2012)

PMID:23178880

el at., Inhibition of DNA-PK enhances chemosensitivity of B-cell precursor acute lymphoblastic leukemia cells to doxorubicin.In Biomed Pharmacother on 2017 Oct by Fatemeh Alikarami, Majid Safa, et al...PMID: 28821159, (2017)

PMID:28821159

Note: This product is for in vitro research use only and is not intended for use in humans or animals.