# **VEGFA Antibody**

Catalog No: #31274

Package Size: #31274-1 50ul #31274-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


# Description

Product Name	VEGFA Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	ELISA WB IHC
Species Reactivity	Human
Specificity	The antibody detects endogenous level of total VEGFA protein.
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide of human VEGFA
Conjugates	Unconjugated
Target Name	VEGFA
Other Names	vascular endothelial growth factor A, VPF, VEGF, MVCD1
Accession No.	P15692
Calculated MW	27 kDa
SDS-PAGE MW	35kd
Concentration	0.8mg/ml
Formulation	Supplied at 0.9mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.3, 0.05% sodium azide
	and 50% glycerol.
Storage	Store at -20°C/1 year

# **Application Details**

WB 1:500-1:2000IHC 1:50-1:200ELISA 1:5000-1:10000

## **Images**




Gel: 8%SDS-PAGE Lysate: 40 ug

Lane: Hela cell lysate

Primary antibody: (VEGFAAntibody) at dilution 1/600 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution

Exposure time: 2 minutes



The image on the left is immunohistochemistry of paraffin-embedded
Human liver cancer tissue using (VEGFA Antibody) at dilution

## Background

This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of, and in-frame with the first AUG, leading to additional isoforms.

#### **Published Papers**

el at., Artesunate Inhibits Proliferation and Invasion of Mouse Hemangioendothelioma Cells in vitro and of Tumor Growth in vivo.In Oncol Lett on 2017 Nov by Ning Wang, Hongxia Chen, et al.. PMID: 29113263, , (2017)

#### PMID:29113263

el at., Photoacoustic molecular imaging-escorted adipose photodynamic-browning synergy for fighting obesity with virus-like complexes. In Nat Nanotechnol on 2021 Apr by Ronghe Chen, Shanshan Huang, et al.. PMID:33526836, (2021)

#### PMID:33526836

el at., Interleukin-17 and vascular endothelial growth factor: new biomarkers for the diagnosis of papillary thyroid carcinoma in patients with Hashimotoβ s thyroiditis. In J Int Med Res on 2022 Jan by Shuiping Li, Shilin Li,et al..PMID: 35023376, , (2022)

#### PMID:35023376

el at., 3D printed PGCL@PLA/10CSPL composite scaffolds loaded with fibronectin 1 for intervertebral disc degeneration treatment, In Biofabrication on 2024 Dec 13 by Weilin Zhang, Siyuan Chen, et al.. PMID:39668784, , (2024)

PMID:39668784

Note: This product is for in vitro research use only and is not intended for use in humans or animals.