

NR3C1 Antibody

Catalog No: #32634



Package Size: #32634-1 50ul #32634-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	NR3C1 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	WB,IHC,IF
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous level of total NR3C1 protein.
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant fusion protein of human Glucocorticoid Receptor (NP_001191194.1).
Conjugates	Unconjugated
Target Name	NR3C1
Other Names	GCCR;GCR;GCRST;GR;GRL;NR3C1
Accession No.	Uniprot:P04150GeneID:2908
SDS-PAGE MW	105kDa
Concentration	1.0mg/ml
Formulation	PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Storage	Store at -20°C. Avoid freeze / thaw cycles.


Application Details

WB 1:500 - 1:2000IHC 1:50 - 1:200IF 1:50 - 1:200


Images




Western blot analysis of extracts of various cell lines, using Glucocorticoid Receptor antibody.




Immunohistochemistry of paraffin-embedded rat spleen using Glucocorticoid Receptor Rabbit pAb.



Immunohistochemistry of paraffin-embedded human liver cancer using Glucocorticoid Receptor Rabbit pAb.



Immunofluorescence analysis of NIH/3T3 cells using Glucocorticoid Receptor Rabbit pAb.



Immunofluorescence analysis of PC-12 cells using Glucocorticoid Receptor Rabbit pAb.

Background

This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).

Published Papers

et al., Disrupted development from head to tail: Pervasive effects of postnatal restricted resources on neurobiological, behavioral, and morphometric outcomes. In Front Behav Neurosci on 2022 Aug 5 by Molly H Kent, Joanna C Jacob, et al..PMID: 35990727, , (2022)

[PMID:35990727](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.