# FLAG tag Antibody

Catalog No: #35535

Package Size: #35535-1 50ul #35535-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escri	ntı	Λn
$\boldsymbol{ u}$	COUL	рu	OH

Product Name	FLAG tag Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antigen affinity purification.	
Applications	ELISA WB	
Specificity	The antibody detects transfected proteins contanining FLAG tag.	
Immunogen Type	Peptide	
Immunogen Description	Synthetic peptide: 3*DYKDDDDK conjugated to KLH	
Target Name	FLAG tag	
Other Names	DDDDK epitope tag; DYKDDDDK epitope tag	
Calculated MW	50/40 kDa	
Concentration	1.0 mg/ml	
Formulation	Rabbit IgG in pH7.3 PBS, 0.05% NaN3, 50% Glycerol.	
Storage	Store at -20°C	

# **Application Details**

WB 1:10000-1:50000 ELISA:1:5000-1:10000

# **Images**



Gel: 8%SDS-PAGE Lysate: 0.1/1 ΞΌg

Lane 1-2: 0.1/1 = Og Recombinant proteins containing the

Flag Tag

Primary antibody: (FLAG Tag Antibody) at dilution 1/10000 Secondary antibody: (HRP-conjugated Goat anti rabbit IgG) at

1/20000 dilution

Exposure time: 30 seconds (Detected by Chemiluminescence

Imager)

# Background

The DYKDDDDK peptide has been used extensively as a general epitope tag in expression vectors. Due to its small size, it is unlikely to affect the tagged protein's biochemical properties. This antibody recognizes the DYKDDDDK peptide fused to either the amino- or carboxy-terminus of targeted proteins. (The background information is obtained from official databases and does not represent the product information.)

# **Published Papers**

TaoChen, Jingjie Li et al, el at., PKC0 phosphorylates MIIP and promotes colorectal cancer metastasis through inhibition of RelA deacetylation., Nature Communications, 8(1):939. doi: 10.1038/s41467-017-01024-2.(2017 Oct 16)

### PMID:29038521

Ting Wangι ε- iujing Yu Li et al el at., GlcNAcylation of fumarase maintains tumour growth under glucose deficiency., Nature Cell Biology, doi: 10.1038/ncb3562(2017)

## PMID:28628081

el at., PAK4 Phosphorylates Fumarase and Blocks TGFβ-Induced Cell Growth Arrest in Lung Cancer Cells. In Cancer Res on 2019 Apr 1 by Chen T, Wang T, et al..PMID:30683654, , (2019)

#### PMID:30683654

el at., MLL2 promotes cancer cell lymph node metastasis by interacting with RelA and facilitating STC1 transcription. In Cell Signal on 2020 Jan; by Li H, Li Q, et al.. PMID:31676369, (2020)

### PMID:31676369

el at., O-GlcNAcylation of PFKFB3 is required for tumor cell proliferation under hypoxia. In Oncogenesis on 2020 Feb 14 by Lei Y, Chen T, et al.. PMID: 32060258, (2020)

### PMID:32060258

el at., Upregulation of ZNF148 in SDHB-deficient gastrointestinal stromal tumor potentiates Forkhead box M1-mediated transcription and promotes tumor cell invasion.

In Cancer Sci on 2020 Apr by Gao X, Ma C, et al..PMID: 32060966, , (2020)

### PMID:32060966

el at., O-GlcNAcylation of Fumarase Maintains Tumour Growth Under Glucose Deficiency .In Nat Cell Biol ON 2017 Jul by Ting Wang, Qiujing Yu,et al..PMID: 28628081, , (2017)

# PMID:28628081

el at., PKC0 phosphorylates MIIP and promotes colorectal cancer metastasis through inhibition of RelA deacetylation. In Nat Commun on 2017 Oct 16 by Tao Chen, Jingjie Li ,et al.. PMID: 29038521, , (2017)

## PMID:29038521

el at., Interaction between NS1 and Cellular MAVS Contributes to NS1 Mitochondria Targeting. In Viruses on 2021 Sep 23 by Yeu-Yang Tseng, Chih-Ying Kuan,et al..PMID:34696339, , (2021)

PMID:34696339

Note: This product is for in vitro research use only and is not intended for use in humans or animals.