

BCL2A1 Antibody

Catalog No: #35596

Package Size: #35596-1 50ul #35596-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


Product Name	BCL2A1 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification.
Applications	WB IHC
Species Reactivity	Human;Mouse
Specificity	The antibody detects endogenous levels of total BCL2A1 protein.
Immunogen Type	Recombinant Protein
Immunogen Description	Fusion protein corresponding to a region derived from internal residues of human BCL2-related protein A1
Conjugates	Unconjugated
Target Name	BCL2A1
Other Names	GRS; BFL1; ACC-1; ACC-2; HBPA1; BCL2L5
Accession No.	Swiss-Prot#: Q16548NCBI Gene ID: 597Gene Accssion: BC016281
SDS-PAGE MW	20kd
Concentration	0.7mg/ml
Formulation	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Storage	Store at -20°C

Application Details

Western blotting: 1:200-1:1000

Immunohistochemistry: 1:25-1:100

Images



Gel: 12%SDS-PAGE


Lysates (from left to right): Mouse testis tissue and hela cell

Amount of lysate: 40ug per lane

Primary antibody: 1/117 dilution

Secondary antibody dilution: 1/8000

Exposure time: 1 minute



Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using #35596 at dilution 1/20.

Background

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is up-regulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Published Papers

el at., KCNN4 induces multiple chemoresistance in breast cancer by regulating BCL2A1. In Am J Cancer Res on 2020 Oct 1 by Peiyang Lin, Junjing Li,et al..PMID:33163271, , (2020)

[PMID:33163271](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.