

FGF2 Antibody

Catalog No: #36769

Package Size: #36769-1 50ul #36769-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


Product Name	FGF2 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification.
Applications	WB IHC
Species Reactivity	Human;Mouse;Rat
Specificity	The antibody detects endogenous levels of total FGF2 protein.
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide corresponding to a region derived from internal residues of human Fibroblast growth factor 2
Conjugates	Unconjugated
Target Name	FGF2
Other Names	BFGF; FGFB; FGF-2; HBGF-2
Accession No.	Swiss-Prot#: P09038NCBI Gene ID: 2247Gene Accssion: NP_001997
SDS-PAGE MW	31kd
Concentration	2.7mg/ml
Formulation	Rabbit IgG in pH7.3 PBS, 0.05% NaN ₃ , 50% Glycerol.
Storage	Store at -20°C

Application Details

Western blotting: 1:1000-1:5000

Immunohistochemistry: 1:100-1:500

Images



Gel: 8%SDS-PAGE


Lysates (from left to right): Mouse lung tissue

Amount of lysate: 30ug per lane

Primary antibody: 1/1000 dilution

Secondary antibody dilution: 1/8000

Exposure time: 10 seconds



Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue using #36769 at dilution 1/100.

Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.

Published Papers

el at., Growth of MCF-7 breast cancer cells and efficacy of anti-angiogenic agents in a hydroxyethyl chitosan/glycidyl methacrylate hydrogel. In Cancer Cell Int on 2017 May 16 by Hejing Wang,Junmin Qian,et al..PMID: 28515673, , (2017)

[PMID:28515673](#)

el at., Apurinic/apurimidinic endonuclease 1 induced upregulation of fibroblast growth factor 2 and its receptor 3 induces angiogenesis in human osteosarcoma cells. In Cancer Sci on 2014 Feb by Tao Ren, Yi Qing et al..PMID:24329908 , , (2014)

[PMID:24329908](#)

el at., ATF6 aggravates angiogenesis-osteogenesis coupling during ankylosing spondylitis by mediating FGF2 expression in chondrocytes. In iScience on 2021 Jun 28 by Mengjun Ma, Hongyu Li,et al..PMID:34296071, , (2021)

[PMID:34296071](#)

Wu Yang;Weiwen Zhu;Yunfei Yang;Minkang Guo;Husun Qian;Weiqian Jiang;Yu Chen;Chengjie Lian;Zijie Xu;Haobo Bai;Tingmei Chen;Jian Zhang el at., Exosomal miR-100-5p inhibits osteogenesis of hBMSCs and angiogenesis of HUVECs by suppressing the BMPR2/Smad1/5/9 signalling pathway., (2021)

[PMID:34256859](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.