CD68 Polyclonal Antibody

Catalog No: #41618

Package Size: #41618-1 50ul #41618-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| _ | | | | |
|---|----|-----|-----|-----|
| П | 00 | ori | nti | ion |
| U | ヒ٥ | UH | บแ | UI |

| Product Name | CD68 Polyclonal Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific |
| | immunogen. |
| Applications | WB ELISA IHC-p IF |
| Species Reactivity | Hu |
| Specificity | CD68 Polyclonal Antibody detects endogenous levels of CD68 protein. |
| Immunogen Description | Synthesized peptide derived from the Internal region of human CD68. |
| Target Name | CD68 |
| Other Names | CD68; Macrosialin; Gp110; CD68 |
| Accession No. | Swiss-Prot: P34810NCBI Gene ID: 968 |
| SDS-PAGE MW | 37kd |
| Concentration | 1mg/ml |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | Store at -20°C/1 year |

Application Details


Western Blot: 1/500 - 1/2000.

ELISA: 1/20000.

Not yet tested in other applications.

IHC-p: 1:100-300

Images



Western Blot analysis of mouse spleen cells using CD68 Polyclonal Antibody

| Note: This product is for in vitro research use only and is not intended for use in humans or animals. | | | | | | |
|--|--|--|--|--|--|--|
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |