

DDC Antibody

Catalog No: #43658

Package Size: #43658-1 50ul #43658-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	DDC Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification
Applications	WB
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total DDC protein.
Immunogen Type	peptide
Immunogen Description	Synthetic peptide of human DDC
Target Name	DDC
Other Names	AADC
Accession No.	Swiss-Prot#: P20711NCBI Gene ID: 1644
Calculated MW	54kd
Concentration	0.7mg/ml
Formulation	Rabbit IgG in pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol.
Storage	Store at -20°C

Application Details

Western blotting: 1:500-2000

Images



Gel: 8%SDS-PAGE

Lysate: 40 µg, Lane 1-2: Human left kidney tissue and Human fetal liver tissue lysates,

Primary antibody:DDC antibody at dilution 1/800,

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution,

Exposure time: 10 seconds

Background

The encoded protein catalyzes the decarboxylation of L-3,4-dihydroxyphenylalanine (DOPA) to dopamine, L-5-hydroxytryptophan to serotonin and L-tryptophan to tryptamine. Defects in this gene are the cause of aromatic L-amino-acid decarboxylase deficiency (AADCD). AADCD deficiency is an inborn error in neurotransmitter metabolism that leads to combined serotonin and catecholamine deficiency. Multiple alternatively spliced transcript

variants encoding different isoforms have been identified for this gene.

Published Papers

el at., 5-HT_{2A} Receptor and 5-HT Degradation Play a Crucial Role in Atherosclerosis by Modulating Macrophage Foam Cell Formation, Vascular Endothelial Cell Inflammation, and Hepatic Steatosis. In J Atheroscler Thromb on 2021 Feb 2 by Yingying Ma, Xiurui Liang,et al..PMID: 33536397, , (2021)

[PMID:33536397](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.