Product Datasheet

p38 Rabbit mAb

Catalog No: #48644

Package Size: #48644-1 50ul #48644-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description	
Product Name	

Product Name	p38 Rabbit mAb	
Host Species	Recombinant Rabbit	
Clonality	Monoclonal antibody	
Clone No.	SR43-04	
Purification	ProA affinity purified	
Applications	WB, ICC/IF,IHC	
Species Reactivity	Human;Mouse;Rat	
Immunogen Description	recombinant protein	
Conjugates	Unconjugated	
Other Names	CSAID Binding Protein 1 antibody CSAID binding protein antibody CSAID-binding protein antibody Csaids	
	binding protein antibody CSBP 1 antibody CSBP 2 antibody CSBP antibody CSBP1 antibody CSBP2 antibody	
	CSPB1 antibody Cytokine suppressive anti-inflammatory drug-binding protein antibody EXIP antibody MAP	
	kinase 14 antibody MAP kinase MXI2 antibody MAP kinase p38 alpha antibody MAPK 14 antibody MAPK14	
	antibody MAX interacting protein 2 antibody MAX-interacting protein 2 antibody Mitogen Activated Protein	
	Kinase 14 antibody Mitogen activated protein kinase p38 alpha antibody Mitogen-activated protein kinase 14	
	antibody Mitogen-activated protein kinase p38 alpha antibody MK14_HUMAN antibody Mxi 2 antibody MXI2	
	antibody p38 ALPHA antibody p38 antibody p38 MAP kinase antibody p38 MAPK antibody p38 mitogen	
	activated protein kinase antibody p38ALPHA antibody p38alpha Exip antibody PRKM14 antibody PRKM15	
	antibody RK antibody SAPK2A antibody	
Accession No.	Swiss-Prot#:Q16539	
Calculated MW	41 kDa	
SDS-PAGE MW	40 kDa	
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.	
Storage	Store at -20°C	


Application Details

WB: 1:1,000-5,000ICC: 1:50-1:200


Images



Western blot analysis of p38 on different cell lysates using anti-p38 antibody at 1/1,000 dilution. Positive control: Lane 1: Hela Lane 2: NIH/3T3 Lane 3: PC-12 Lane 4: Jurkat



ICC staining p38 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining p38 in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

MAP (mitogen-activated protein) kinases play a significant role in many biological processes, including cell adhesion and spreading, cell differentiation and apoptosis. $p38\alpha$, $p38\beta$ and $p38\gamma$, also known as MAPK14, MAPK11 and MAPK12, respectively, each contain one protein kinase domain and belong to the MAP kinase family. Expressed in different areas throughout the body with common expression patterns in heart, p38 proteins use magnesium as a cofactor to catalyze the ATP-dependent phosphorylation of target proteins. Via their catalytic activity, $p38\alpha$, $p38\beta$ and $p38\gamma$ are involved in a variety of events throughout the cell, including signal transduction pathways, cytokine production and cell proliferation and differentiation. The p38 proteins are subject to phosphoryation on Thr and Tyr residues, an event which is thought to activate the phosphorylated protein.

References

1. Yan T et al. Luteolin inhibits behavioral sensitization by blocking methamphetamine-induced MAPK pathway activation in the caudate putamen in mice. PLoS One 9:e98981 (2014). 2. Pagliara V et al. Protease Nexin-1 affects the migration and invasion of C6 glioma cells through the regulation of urokinase Plasminogen Activator and Matrix Metalloproteinase-9/2. Biochim Biophys Acta 1843:2631-44 (2014).

Published Papers

el at., Confirming whether novel rhein derivative 4a induces paraptosis-like cell death by endoplasmic reticulum stress in ovarian cancer cells. In Eur J Pharmacol on 2020 Nov 5 by Hui-Feng Pang, Xin-Xiao Li, et al..PMID:32890460, , (2020)

PMID:32890460

el at., GFAT1-linked TAB1 glutamylation sustains p38 MAPK activation and promotes lung cancer cell survival under glucose starvation. In Cell Discov on 2022 Aug 9 by Shupei Wei, Qin Zhao, et al..PMID:35945223, , (2022)

PMID:35945223

el at., Novel oxicam nonsteroidal compound XK01 attenuates inflammation by suppressing the NF-kB and MAPK pathway in RAW264.7 macrophages. In Heliyon on 2024 Jan 9 by Jixiang Wang, Jiawang Tan,et al..PMID:38312593, , (2024)

PMID:38312593

Note: This product is for in vitro research use only and is not intended for use in humans or animals.