

## CD8 alpha Rabbit mAb

Catalog No: #48750



Package Size: #48750-1 50ul #48750-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	CD8 alpha Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SI18-01
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, FC
Species Reactivity	Human
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	alpha polypeptide (p32) antibody CD8 antibody CD8 antigen alpha polypeptide antibody CD8 antigen alpha polypeptide (p32) antibody CD8a antibody CD8A antigen antibody CD8A molecule antibody CD8A_HUMAN antibody Leu2 antibody Leu2 T lymphocyte antigen antibody Ly3 antibody LYT3 antibody MAL antibody OKT8 T cell antigen antibody OTTHUMP00000160760 antibody OTTHUMP00000160764 antibody OTTHUMP00000203528 antibody OTTHUMP00000203721 antibody p32 antibody T cell antigen Leu2 antibody T cell co receptor antibody T-cell surface glycoprotein CD8 alpha chain antibody T-lymphocyte differentiation antigen T8/Leu-2 antibody T8 T cell antigen antibody T8/Leu-2 T-lymphocyte differentiation antigen antibody
Accession No.	Swiss-Prot#:P01732
Calculated MW	26 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200 FC: 1:500-1:1,000


## Images




Western blot analysis of CD8 alpha on different lysates using anti-CD8 alpha antibody at 1/1,000 dilution. Positive control:  
Lane 1: HeLa      Lane 2: Lovo



Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-CD8 alpha antibody. Counter stained with hematoxylin.



ICC staining CD8 alpha in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CD8 alpha in CRC cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

## Background

The T cell receptor (TCR) is a heterodimer composed of either  $\alpha$  and  $\beta$  or  $\gamma$  and  $\delta$  chains. CD3 chains and the CD4 or CD8 (CD8- $\alpha$  and CD8- $\beta$ ) co-receptors are also required for efficient signal transduction through the TCR. The TCR is expressed on T helper and T cytotoxic cells that can be distinguished by their expression of CD4 and CD8 proteins; T helper cells express CD4 proteins and T cytotoxic cells display CD8 proteins. CD8s are cell surface glycoproteins that exist as two chain complex ( $\alpha\alpha$  or  $\alpha\beta$ ) receptors that bind class I MHC molecules presented by the antigen-presenting cell (APC). A primary function of CD8 proteins is to facilitate antigen recognition by the TCR and to strengthen the avidity of the TCR-antigen interactions. An additional role for CD8-expressing T cells may be to maintain low levels of HIV expression.

## References

1. Zehbe, I. et al. 2007. Human papillomavirus 16 E6-specific CD45RA+ CCR7+ high avidity CD8+ T cells fail to control tumor growth despite interferon-gamma production in patients with cervical cancer. *J Immunother.* 30: 523-532.
2. Morales, P.J. et al. 2003. Placental cell expression of HLA-G2 isoforms is limited to the invasive trophoblast phenotype. *J. Immunology.* 171: 6215-6224.

## Published Papers

el at., DKK1 affects survival of patients with head and neck squamous cell carcinoma by inducing resistance to radiotherapy and immunotherapy  
*In Radiother Oncol* 2023 Apr by Xinyu Ye?1, Jingwen Liu? et al.. PMID: 36690301, (2023)

[PMID:36690301](#)

---

Zhao YuLong, Li YouJie, Wu ChunXia, Xie Ning, Li XiYan, Cao DianChao, Guo Yang, Liang Yan, Li YanMei, Xue JiangNan, Sun HongFang, Wang Qin, Li XiaoHua, Wang PingYu, Xie ShuYang et al., m6A-modified circZNF548 regulates exosomal miR-7108-3p to activate CD3+CD8+ T cells and suppress NSCLC growth by JMY, BMC biology, (2025)

[PMID:40817234](#)

---

Zhang Meiling, Xie Ning, Wu Dongyang, Wu Chunxia, Guo Yang, Wang Na-Na, Li Xi-Yan, Liang Yan, Xie Xinghan, Wang Tengfei, Li You-Jie, Ning Fangling, Wang Ranran, Xie Shu-Yang, Wang Pingyu et al., Novel nanoparticle-combined therapy through EGFR and PD-L1 blockade: PFPR-siRNA-mediated suppressing NSCLC proliferation via PDT and enhanced antitumor  $\beta$   $\beta$ , Colloids and surfaces. B, Biointerfaces, (2025)

[PMID:41265187](#)

---

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.