# Myosin light chain kinase Rabbit mAb

Catalog No: #48846

Package Size: #48846-1 50ul #48846-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_		4.5	
Des	cri	ntic	าท
レしる	UH	$\rho$ u	ווע

Product Name	Myosin light chain kinase Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SU40-06
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, FC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	deglutamylated form antibody DKFZp686l10125 antibody EC 2.7.11.18 antibody FLJ12216 antibody Kinase
	related protein antibody Kinase-related protein antibody KRP antibody MLCK antibody MLCK1 antibody
	MLCK108 antibody MLCK210 antibody MSTP083 antibody MYLK antibody MYLK_HUMAN antibody MYLK1
	antibody Myosin light chain kinase antibody Myosin light polypeptide kinase antibody OTTHUMP00000180642
	antibody OTTHUMP00000180643 antibody smMLCK antibody smooth muscle antibody Smooth muscle
	myosin light chain kinase antibody Telokin antibody
Accession No.	Swiss-Prot#:Q15746
Calculated MW	210 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C
Storage	Store at -20°C


## Application Details

WB: 1:2,000-1:10,000 IHC: 1:100-1:500 ICC: 1:100-1:500 FC: 1:50-1:100


## **Images**




Western blot analysis of MYLK on human lung lysates using anti-MYLK antibody at 1/5,000 dilution.




Immunohistochemical analysis of paraffin-embedded rat smooth muscle tissue using anti-MYLK antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue using anti-MYLK antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse smooth muscle tissue using anti-MYLK antibody. Counter stained with hematoxylin.




ICC staining MYLK in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




ICC staining MYLK in L6 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining MYLK in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining MYLK in SH-SY-5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of SH-SY-5Y cells with MYLK antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

#### Background

MLCK, a member of the Ser/Thr protein kinase family, is a calcium/calmodulin-dependent enzyme responsible for smooth muscle contraction via phosphorylation of a specific serine in the N-terminus of myosin light chains (MLC), an event that facilitates myosin interaction with actin filaments. It is a central determinant in the development of vascular permeability and tissue edema formation. In the nervous system it has been shown to control the growth initiation of astrocytic processes in culture and to participate in transmitter release at synapses formed between cultured sympathetic ganglion cells. MLCK acts as a critical participant in signaling sequences that result in fibroblast apoptosis. Smooth muscle and non-muscle isozymes are expressed in a wide variety of adult and fetal tissues and in cultured endothelium with qualitative expression appearing to be neither tissue- nor development-specific. Non-muscle isoform 2 is the dominant splice variant expressed in various tissues. The Telokin isoform, which binds calmodulin, has been found in a wide variety of adult and fetal tissues. MLCK is probably down-regulated by phosphorylation. The protein contains 1 fibronectin type III domain and 9 immunoglobulin-like C2-type domains.

### References

- 1. Liang QX et al. Deletion of Mylk1 in oocytes causes delayed morula-to-blastocyst transition and reduced fertility without affecting folliculogenesis and oocyte maturation in mice. Biol Reprod 92:97 (2015).
- 2. Liu FF et al. Characteristics of diprophylline-induced bidirectional modulation on rat jejunal contractility. Korean J Physiol Pharmacol 18:47-53 (2014).

## **Published Papers**

el at., Gastrodin attenuates angiotensin II-induced vascular contraction and MLCK/p-MLC2 pathway activationIn Pharm BiolOn2023 DecbyZhi Guo , Xuan Yang et al..PMID:37211627, , (2023)

PMID:37211627

Note: This product is for in vitro research use only and is not intended for use in humans or animals.