

MUC1 Rabbit mAb

Catalog No: #49024



Package Size: #49024-1 50ul #49024-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com






Description



Product Name	MUC1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SN06-80
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Other Names	ADMCKD antibody ADMCKD1 antibody Breast carcinoma associated antigen DF3 antibody Breast carcinoma-associated antigen DF3 antibody CA 15-3 antibody CA15 3 antibody CA15 3 antigen antibody CA15.3 antibody Cancer antigen 15-3 antibody Carcinoma associated mucin antibody Carcinoma-associated mucin antibody CD 227 antibody CD227 antibody DF3 antigen antibody EMA antibody Episialin antibody H23 antigen antibody H23AG antibody KL 6 antibody KL-6 antibody KL6 antibody Krebs von den Lungen-6 antibody MAM 6 antibody MAM6 antibody MCD antibody MCKD antibody MCKD1 antibody Medullary cystic kidney disease 1 (autosomal dominant) antibody Medullary cystic kidney disease, autosomal dominant antibody MUC 1 antibody MUC-1 antibody MUC-1/SEC antibody MUC-1/X antibody MUC1 antibody MUC1-alpha antibody MUC1-beta antibody MUC1-CT antibody MUC1-NT antibody MUC1/ZD antibody MUC1_HUMAN antibody Mucin 1 antibody Mucin 1 transmembrane antibody Mucin 1, cell surface associated antibody Mucin-1 subunit beta antibody Peanut reactive urinary mucin antibody Peanut-reactive urinary mucin antibody PEM antibody PEMT antibody Polymorphic epithelial mucin antibody PUM antibody Tumor associated epithelial membrane antigen antibody Tumor associated epithelial mucin antibody Tumor associated mucin antibody Tumor-associated epithelial membrane antigen antibody Tumor-associated mucin antibody
Accession No.	Swiss-Prot#:P15941
Calculated MW	23 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100

Images





Background

The mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs) encoded by 60 base pairs (Mucin 1), 69 base pairs (Mucin 2) and 51 base pairs (Mucin 3). The number of repeats is highly polymorphic and varies among different alleles. Mucin 1 proteins are expressed as type I membrane proteins in addition to secreted forms. Mucin 1 is aberrantly expressed in epithelial tumors including breast carcinomas. Mucin 2 coats the epithelia of the intestines and airways and is associated with colonic tumors. Mucin 3 is a major component of various mucus gels and is broadly expressed in normal and tumor cells.

References

1. Gorges TM et al. Circulating tumour cells escape from EpCAM-based detection due to epithelial-to-mesenchymal transition. BMC Cancer 12:178 (2012).
2. Williams MA et al. Deletion of the mucin-like molecule muc1 enhances dendritic cell activation in response to toll-like receptor ligands. J Innate Immun 2:123-43 (2010).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.