

## NQO1 Rabbit mAb

Catalog No: #49364



Package Size: #49364-1 50ul #49364-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)


## Description

Product Name	NQO1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JF440-1
Purification	ProA affinity purified
Applications	WB, ICC/IF, IP, FC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Azoreductase antibody Cytochrome b 5 reductase antibody DHQU antibody DIA 4 antibody DIA4 antibody Diaphorase (NADH/NADPH) (cytochrome b 5 reductase) antibody Diaphorase (NADH/NADPH) antibody Diaphorase 4 antibody Dioxin inducible 1 antibody DT diaphorase antibody DT-diaphorase antibody DTD antibody Menadione reductase antibody NAD(P)H dehydrogenase [quinone] 1 antibody NAD(P)H dehydrogenase quinone 1 antibody NAD(P)H menadione oxidoreductase 1 dioxin inducible antibody NAD(P)H: menadione oxidoreductase 1 dioxin inducible 1 antibody NAD(P)H:menadione oxidoreductase 1 antibody NAD(P)H:Quinone acceptor oxidoreductase type 1 antibody NAD(P)H:quinone oxidoreductase 1 antibody NAD(P)H:quinone oxireductase antibody NMOR 1 antibody NMOR I antibody NMOR1 antibody NMORI antibody NQO 1 antibody NQO1 antibody NQO1_HUMAN antibody Phylloquinone reductase antibody QR 1 antibody QR1 antibody Quinone reductase 1 antibody
Accession No.	Swiss-Prot#:P15559
Calculated MW	31/27 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:1,000-5,000 ICC: 1:50-1:200 IP: 1:50 FC: 1:50-1:100

## Images



Western blot analysis of NQO1 on different lysates using anti-NQO1 antibody at 1/1,000 dilution. Positive control:  
Lane 1: SH-SY-5Y Lane 2: Mouse kidney Lane 3: Raji



Flow cytometric analysis of HeLa cells with NQO1 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody

## Background

NAD(P)H:quinone oxidoreductase 1 (NQO1) and NRH:quinone oxidoreductase (NQO2) are flavoproteins that catalyze the metabolic detoxification of quinones and their derivatives to hydroquinones, using either NADH or NADPH as the electron donor. This protects cells against quinone-induced oxidative stress, cytotoxicity, and mutagenicity. Many tumors overexpress NQO1, which is an obligate two-electron reductase that deactivates toxins and activates bioreductive anticancer drugs. NQO1, a 274 amino acid protein, is ubiquitously expressed, but the expression level varies among tissues. NQO1 gene expression is coordinately induced in response to xenobiotics, antioxidants, heavy metals and radiation. The antioxidant response element (ARE) in the NQO1 gene promoter is essential for expression and coordinated induction of NQO1. ARE activation by tert-butylhydroquinone is dependent on PI3-kinase, which lies upstream of Nrf2. Nrf2, c-Jun, Nrf1, Jun-B and Jun-D bind to the ARE and regulate expression and induction of NQO1 gene. Maf-Maf homodimers and possibly Maf-Nrf2 heterodimers play a role in negative regulation of ARE-mediated transcription, but Maf-Nrf1 heterodimers fail to bind with the NQO1 gene ARE and do not repress NQO1 transcription.

## References

1. Greco T et al. Ketogenic diet decreases oxidative stress and improves mitochondrial respiratory complex activity. *J Cereb Blood Flow Metab* 36:1603-13 (2016).
2. Kigoshi Y et al. CACUL1/CAC1 Regulates the Antioxidant Response by Stabilizing Nrf2. *Sci Rep* 5:12857 (2015).

## Published Papers

el at., Asperuloside suppressing oxidative stress and inflammation in DSS-induced chronic colitis and RAW 264.7 macrophages via Nrf2/HO-1 and NF- $\kappa$ B pathways. In *Chem Biol Interact* on 2021 Aug 1 by Yong-Er Chen , Shi-Jie Xu et al..PMID:33974900, , (2021), , (2021), , (2021)  
[PMID:33974900](#)

Jin Yuan;Junwei Su;Shaowen Zhong;Xin Yuan;Jianping Zhu;Shuangxi Lu;Di Zhang;Guiling Li;Hanyu Xue;Min Yan;Lei Yue;Tianwu Zhang et al., Dictamnine alleviates DSS-induced colitis mice by inhibiting ferroptosis of enterocytes via activating Nrf2-Gpx4 signaling pathway., (2025)  
[PMID:40049578](#)

Duan Hong, Zhang Yaci, Shen Aijuan, Ren Jiahui, Zhang Fengxia, Lu Yunshuo, Wei Xuedan, Yang Chaoyu, Gong Jiexi, Wang Xin, Du Yongkun, Pei Qiming, Zhang Angke et al., Salvianolic acid A inhibits PRRSV replication via binding to Keap1 to activate the MKRN1 $\beta$  Nrf2 $\beta$  NQO1 pathway, *Veterinary research*, (2025)

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.