

DAP12 Rabbit mAb

Catalog No: #49944



Package Size: #49944-1 50ul #49944-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

| | |
|-----------------------|---|
| Product Name | DAP12 Rabbit mAb |
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal antibody |
| Clone No. | JG38-70 |
| Purification | ProA affinity purified |
| Applications | WB,IHC,FC |
| Species Reactivity | Human |
| Immunogen Description | Synthetic peptide within human DAP12 aa 60-100. |
| Conjugates | Unconjugated |
| Other Names | DAP 12 antibody DAP12 antibody DNAX activation protein 12 antibody DNAX-activation protein 12 antibody KAR-associated protein antibody KARAP antibody Killer activating receptor associated protein antibody Killer-activating receptor-associated protein antibody PLOSL antibody TYOBP_HUMAN antibody TYRO protein tyrosine kinase binding protein antibody TYRO protein tyrosine kinase-binding protein antibody TYROBP antibody |
| Accession No. | Swiss-Prot#:O43914 |
| Calculated MW | 12 kDa |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |


Application Details

WB: 1:500-1:1,000 IHC: 1:50-1:200FC: 1:50-1:100


Images



Western blot analysis of DAP12 on THP-1 cell using anti-DAP12 antibody at 1/500 dilution.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue using anti-DAP12 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of K562 cells with DAP12 antibody at 1/100 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

Background

Natural killer (NK) cells are regulated by stimulatory and inhibitory signals from a variety of receptors. Three main receptor families are responsible for NK cells recognition of MHC I molecules, including Ly-49, CD94/NKG2 and KIR (killer-cell inhibitory receptor). DAP12 is a phosphoprotein that is involved in the activation of NK cells. This protein interacts with membrane glycoproteins of the KIR family, resulting in cellular activation. DAP12 also binds to CD94/NKG2C, an activating NK cell receptor belonging to the C-type lectin superfamily. Additional proteins that bind to DAP12 include Ly-49D and Ly-49H, which associate with DAP12 in the plasma membrane. Phosphorylated DAP12 binds to ZAP-70 and Syk, suggesting that the activation pathway may be similar to that of the T and B cell antigen receptors.

References

1. Paloneva J et al. Loss-of-function mutations in TYROBP (DAP12) result in a presenile dementia with bone cysts. *Nat Genet* 25:357-361 (2000).
2. Kondo T et al. Heterogeneity of presenile dementia with bone cysts (Nasu-Hakola disease): three genetic forms. *Neurology* 59:1105-1107 (2002).

Published Papers

Jingsheng Cai;Zheng Liu;Shaoyi Chen;Jingwei Zhang;Haoran Li;Xun Wang;Feng Yang;Shaodong Wang;Xiao Li;Yun Li;Kezhong Chen;Jun Wang;Ming Sun;Mantang Qiu et al., Engineered circular RNA-based DLL3-targeted CAR-T therapy for small cell lung cancer., , (2025)
[PMID:40075480](https://pubmed.ncbi.nlm.nih.gov/40075480/)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.