CD97 Rabbit mAb

Catalog No: #56568

Package Size: #56568-1 50ul #56568-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| _ | | | | |
|------------------|------|----|-----|-----|
| | esci | 'n | tıc | ۱n |
| \boldsymbol{L} | しつい | ıv | นบ | , , |


| Product Name | CD97 Rabbit mAb |
|-----------------------|--|
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB IHC |
| Species Reactivity | Human;Rat |
| Specificity | CD97 Antibody detects endogenous levels of total CD97 |
| Immunogen Description | A synthesized peptide derived from human CD97 |
| Conjugates | Unconjugated |
| Other Names | ADGRE5; CD97; TM7LN1; |
| Accession No. | Uniprot:P48960 |
| Calculated MW | 92kDa |
| SDS-PAGE MW | 28,55kDa |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |
| | |

Application Details

WB:1:500~1:2000 IHC:1:50~1:200

FC:1:50

Images



Western blot analysis of CD97 expression in human tonsil cell lysate.

Product Description

Receptor potentially involved in both adhesion and signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.

| Note: This product is for in vitro research use only and is not intended for use in humans or animals. | | | | | |
|--|--|--|--|--|--|
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |