

HRPT2 / CDC73 Rabbit mAb

Catalog No: #57154



Package Size: #57154-1 50ul #57154-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	HRPT2 / CDC73 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IHC
Species Reactivity	Human;Mouse;Rat
Specificity	HRPT2 / CDC73 Antibody detects endogenous levels of total HRPT2 / CDC73
Immunogen Description	A synthesized peptide derived from human HRPT2 / CDC73
Conjugates	Unconjugated
Other Names	Cdc73; FIHP; HPTJT; HRPT1; HRPT2; Hyperparathyroidism 2; HYX;
Accession No.	Q6P1J9
Calculated MW	61kDa
SDS-PAGE MW	61kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

WB 1:500~1:2000

Images



All lanes: HRPT2 / CDC73 Rabbit mAb at 1/1k dilution

Lane 1 : 293 whole cell lysates Lane 2 : MCF-7 whole cell lysates Lane 3 : K562 whole cell lysates
Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 61 kDa

Observed band size: 61 kDa

Exposure time: 5 seconds



All lanes: HRPT2 / CDC73 Rabbit mAb at 1/1k dilution

Lane 1 : Mouse lung lysates Lane 2 : Mouse kidney lysates
Lane 3 : Mouse brain lysates Lane 4 : Rat brain lysates Lane 5 : Rat heart lysates
Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 61 kDa

Observed band size: 61 kDa


Exposure time: 5 seconds



All lanes:HRPT2 / CDC73 Rabbit mAb at 1/1k dilution

Lane 1 : Wild-type HeLa cell lysate
Lane 2 : HRPT2 / CDC73 knockdown HeLa cell lysate

Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human kidney tissue stained for HRPT2 / CDC73 using 57154 at 1/100 dilution in immunohistochemical analysis.

Background

Tumor suppressor probably involved in transcriptional and post-transcriptional control pathways. May be involved in cell cycle progression through the regulation of cyclin D1/PRAD1 expression.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.