

Hsc70 Rabbit mAb

Catalog No: #58784



Package Size: #58784-1 50ul #58784-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Hsc70 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	Hsc70 Antibody detects endogenous levels of total Hsc70
Conjugates	Unconjugated
Other Names	Heat shock cognate 71 kDa protein; HS7C; HSP73; HSP7C; HSPA10; HSPA8;
Accession No.	P11142
Calculated MW	71 kDa
SDS-PAGE MW	71 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Application Details

WB: 1:500-1:2000


IHC: 1:50-1:200

ICC/IF: 1:50-1:200


Images




All lanes: Hsc70 Rabbit mAb at 1/1k dilution
 Lane 1 : HeLa whole cell lysates
 Lane 2 : C6 whole cell lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 71 kDa
 Observed band size: 71 kDa
 Exposure time: 4 seconds




All lanes: Hsc70 Rabbit mAb at 1/1k dilution
 Lane 1 : Mouse brain lysates
 Lane 2 : Rat brain lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 71 kDa
 Observed band size: 71 kDa
 Exposure time: 3 seconds



All lanes: Hsc70 Rabbit mAb at 1/1k dilution
 Lane 1 : Wild-type HeLa cell lysate
 Lane 2 : Hsc70 knockdown HeLa cell lysate
 Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human breast cancer tissue stained for Hsc70 using 58784 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence Hsc70 antibody (58784)
 ICC/IF staining of Hsc70 in MCF7 cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 58784 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.