

HDAC3 Rabbit mAb

Catalog No: #58888



Package Size: #58888-1 50ul #58888-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	HDAC3 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	HDAC3 Antibody detects endogenous levels of HDAC3
Conjugates	Unconjugated
Other Names	HD3; Histone deacetylase 3; RPD3-2; SMAP45; HDAC3;
Accession No.	O15379
Calculated MW	49 kDa
SDS-PAGE MW	49 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Application Details

WB: 1:500-1:2000


IHC: 1:50-1:200

ICC/IF: 1:50-1:200


Images




All lanes: HDAC3 Rabbit mAb at 1/1k dilution
 Lane 1 : 293 whole cell lysates
 Lane 2 : MCF-7 whole cell lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 49 kDa
 Observed band size: 49 kDa
 Exposure time: 5 seconds




All lanes: HDAC3 Rabbit mAb at 1/1k dilution
 Lane 1 : RAW264.7 whole cell lysates
 Lane 2 : 3T3 whole cell lysates
 Lane 3 : PC-12 whole cell lysates
 Lane 4 : C6 whole cell lysates
 Lysates/proteins at 20 µg per lane.
 Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 49 kDa
 Observed band size: 49 kDa
 Exposure time: 6 seconds



All lanes : HDAC3 Rabbit mAb at 1/1k dilution
 Lane 1 : Wild-type HAP1 cell lysate
 Lane 2 : HDAC3 knockdown HAP1 cell lysate
 Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human ovarian carcinoma tissue stained for HDAC3 using 58888 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence HDAC3 antibody (58888)
 ICC/IF staining of HDAC3 in HeLa cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 58888 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 647 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.