

## hnRNP K Rabbit mAb

Catalog No: #58915



Package Size: #58915-1 50ul #58915-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

Product Name	hnRNP K Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	hnRNP K Antibody detects endogenous levels of hnRNP K
Conjugates	Unconjugated
Other Names	65 kDa phosphoprotein; CSBP; TUNP; HNRNPK; HNRPK; DC-stretch binding protein; hnRNP-K;
Accession No.	P61978
Calculated MW	51 kDa
SDS-PAGE MW	60 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


## Application Details

WB: 1:500-1:2000


IHC: 1:50-1:200

ICC/IF: 1:50-1:200


## Images




All lanes: hnRNP K Rabbit mAb at 1/1k dilution  
 Lane 1 : Hela whole cell lysates  
 Lane 2 : JK whole cell lysates  
 Lane 3 : HepG2 whole cell lysates  
 Lane 4 : MCF-7 whole cell lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution  
 Predicted band size: 51 kDa  
 Observed band size: 60 kDa  
 Exposure time: 5 seconds




All lanes: hnRNP K Rabbit mAb at 1/1k dilution  
 Lane 1 : Mouse brain lysates  
 Lane 2 : Rat brain lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution  
 Predicted band size: 51 kDa  
 Observed band size: 60 kDa  
 Exposure time: 4 seconds



All lanes: hnRNP K Rabbit mAb at 1/1k dilution  
 Lane 1 : Wild-type HeLa cell lysate  
 Lane 2 : hnRNP K knockdown HeLa cell lysate  
 Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human colon tissue stained for hnRNP K using 58915 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence hnRNP K antibody (58915) ICC/IF staining of hnRNP K in MCF7 cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 58915 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.