Cdc23/APC8 Rabbit mAb

Catalog No: #59168

Package Size: #59168-1 50ul #59168-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

Description


Product Name	Cdc23/APC8 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	Cdc23/APC8 Antibody detects endogenous levels of total Cdc23/APC8
Conjugates	Unconjugated
Other Names	CDC23; ANAPC8; APC8; Cell division cycle 23; CUT23; Cyclosome subunit 8;
Accession No.	Q9UJX2
Calculated MW	69 kDa
SDS-PAGE MW	64 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Application Details


WB: 1:500-1:2000

ICC/IF: 1:50-1:200


Images




All lanes: Cdc23/APC8 Rabbit mAb at 1/1k dilutionLane 1 : HepG2 whole cell lysates Lane 2 : JK whole cell lysates Lysates/proteins at 20 µg per lane.SecondaryAll lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilutionPredicted band size: 69 kDa Observed band size: 64 kDaExposure time: 5 seconds



All lanes: Cdc23/APC8 Rabbit mAb at 1/1k dilutionLane 1 : Mouse spleen lysates Lane 2 : Mouse lung lysates Lane 3 : Rat kidney lysates Lane 4 : Rat brain lysates Lysates/proteins at 20 µg per lane.SecondaryAll lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilutionPredicted band size: 69 kDa Observed band size: 64 kDaExposure time: 7 seconds



All lanes:Cdc23/APC8 Rabbit mAb at 1/1k dilutionLane 1 : Wild-type Hela cell lysateLane 2 :Cdc23/APC8 Rabbit mAb knockdown Hela cell lysateLysates/proteins at 20 µg per lane.



Immunocytochemistry/ Immunofluorescence Cdc23/APC8 antibody (59168) ICC/IF staining of Cdc23/APC8 in Hela cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 59168 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.