

Src Rabbit mAb

Catalog No: #59230



Package Size: #59230-1 50ul #59230-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Src Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	Src Antibody detects endogenous levels of total Src
Conjugates	Unconjugated
Other Names	EC 2.7.10.2; p60-Src; c-Src; pp60;c-src;
Accession No.	P12931
Calculated MW	60 kDa
SDS-PAGE MW	60 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Application Details

WB: 1:500-1:2000


IHC: 1:50-1:200

ICC/IF: 1:50-1:200


Images




All lanes: Src Rabbit mAb at 1/1k dilution
 Lane 1 : 293 whole cell lysates
 Lane 2 : A375 whole cell lysates
 Lane 3 : C6 whole cell lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 60 kDa
 Observed band size: 60 kDa
 Exposure time: 7 seconds




All lanes: Src Rabbit mAb at 1/1k dilution
 Lane 1 : Mouse spleen lysates
 Lane 2 : Mouse lung lysates
 Lane 3 : Rat spleen lysates
 Lane 4 : Rat heart lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 60 kDa
 Observed band size: 60 kDa
 Exposure time: 6 seconds



All lanes: Src Rabbit mAb at 1/1k dilution
 Lane 1 : Wild-type A549 cell lysate
 Lane 2 : Src Rabbit mAb knockdown A549 cell lysate
 Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human kidney tissue stained for Src using 59230 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence Src antibody (59230)
 ICC/IF staining of Src in HeLa cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 59230 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.