

## Annexin V Rabbit mAb

Catalog No: #59326



Package Size: #59326-1 50ul #59326-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description


Product Name	Annexin V Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	Annexin V Antibody detects endogenous levels of total Annexin V
Conjugates	Unconjugated
Other Names	Annexin A5; Anchorin CII; Annexin V; Annexin-5 ; Calphobindin I; CBP-I; Endonexin II; Lipocortin V; PP4; PAP-I; Thromboplastin inhibitor; VAC-alpha; ANXA5;
Accession No.	P08758
Calculated MW	36 kDa
SDS-PAGE MW	33 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Application Details


WB: 1:500-1:2000

ICC/IF: 1:50-1:200


## Images




All lanes: Annexin V Rabbit mAb at 1/1k dilution  
 Lane 1 : 293 whole cell lysates  
 Lane 2 : MCF-7 whole cell lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution  
 Predicted band size: 36 kDa  
 Observed band size: 33 kDa  
 Exposure time: 7 seconds



All lanes: Annexin V Rabbit mAb at 1/1k dilution  
 Lane 1 : Rat heart lysates  
 Lane 2 : Rat kidney lysates  
 Lane 3 : Mouse spleen lysates  
 Lane 4 : Mouse lung lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary: All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution  
 Predicted band size: 36 kDa  
 Observed band size: 33 kDa  
 Exposure time: 4 seconds



All lanes: Annexin V Rabbit mAb at 1/1k dilution  
 Lane 1 : Wild-type A549 cell lysate  
 Lane 2 : Annexin V Rabbit mAb knockdown A549 cell lysate  
 Lysates/proteins at 20 µg per lane.



Immunocytochemistry/ Immunofluorescence Annexin V antibody (59326) ICC/IF staining of Annexin V in HeLa cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 59326 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.