

# Mouse Anti-Human CD133, Biotin Conjugated mAb

Catalog No: #C004

Package Size: #C004 25ug

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Mouse Anti-Human CD133, Biotin Conjugated mAb
Host Species	Mouse
Clonality	Monoclonal
Clone No.	6B3
Isotype	Mouse IgG1, $\kappa$
Applications	FC
Species Reactivity	Hu
Specificity	This antibody recognizes human CD133-2 in FACS. It can not cross react with CD133-1.
Immunogen Description	L929/CD133-2 transfected cells
Other Names	AC133
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in phosphate buffered saline (PBS) and reconstitute with sterile PBS
Storage	Store protected from light at 2-8°C. Do not freeze. The expiration date is indicated on the vial label.

## Application Details

**Preparation:** This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a mouse immunized with L929/CD133-2 transfected cells). The monoclonal antibody was purified from tissue culture supernatant or ascites by protein G affinity chromatography.

**Product Notices:** This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1  $10^6$  cells in a 100- $\mu$ l experimental sample (per test). An isotype control should be used at the same concentration as the antibody of interest.

## Product Description

CD133, known as AC133, is a member of a novel family of cell surface proteins that has five transmembrane domain molecules with an extracellular N-terminus and a cytoplasmic C-terminus, and has two large extracellular loops with eight consensus sites for N-linked glycosylation. It has two isoforms. AC133-1 mRNA was prominent in fetal brain and adult skeletal muscle but was not detected in fetal liver and kidney, adult pancreas, kidney, and placenta. AC133-2 mRNA was found to be dominant in a variety of human fetal tissues, adult tissues, and several carcinomas. CD133 has not only been used to isolate hematopoietic stem cells but also represent a marker of tumor initiating cells in a number of human cancers, such as in human pancreatic adenocarcinoma, colon cancer, melanoma, hepatocellular carcinoma, and brain tumors.

**Note:** This product is for in vitro research use only and is not intended for use in humans or animals.